

U.S. Serial No. 08/477,983

APPLICATION FOR UNITED STATES PATENT



#25-
marked-up
copy

Inventor(s): Jeffrey S. Rubin, Paul W. Finch, Stuart A. Aaronson

Invention: [DNA ENCODING A GROWTH FACTOR SPECIFIC FOR
EPITHELIAL CELLS]

A N EPITHELIAL CELL SPECIFIC GROWTH
FACTOR KERATINOCYTE GROWTH FACTOR (KGF)

Cushman, Darby & Cushman
1615 L Street, N.W.
Eleventh Floor
Washington, D.C. 20036-5601
Attorneys
Telephone: 861-3000

SPECIFICATION

This application is a continuation application of application Serial No. 08/106,775, filed August 16, 1993, abandoned, which is a continuation application of Serial No. 07/780,847, filed October 23, 1991, abandoned, which is a continuation application of Serial No. 07/304,281, filed January 31, 1989, abandoned.

FIELD OF THE INVENTION

The present invention relates to growth factors, particularly to isolation of a polypeptide growth factor similar to a family of factors including known fibroblast growth factors (FGFs). This invention also relates to construction of complementary DNA (cDNA) segments from messenger RNA (mRNA) encoding the novel growth factor. Further, this invention pertains to synthesis of products of such DNA segments by recombinant cells, and to the manufacture and use of certain other novel products enabled by the identification and cloning of DNAs encoding this growth factor.

ABBREVIATIONS USED IN THIS APPLICATION

	aFGF	acidic fibroblast growth factor
	bFGF	basic fibroblast growth factor
	EGF	epidermal growth factor
5	HSAC	heparin-Sepharose affinity chromatography
	kb	kilobases
	kDa	kilodaltons
	KGF	keratinocyte growth factor
10	NaDodSO ₄ /PAGE	Sodium dodecylsulphate (SDS)/ polyacrylamide gel electrophoresis
	RP-HPLC	reversed-phase high performance liquid chromatography
	TGF α	transforming growth factor α

BACKGROUND OF THE INVENTION

Growth factors are important mediators of intercellular communication. These potent molecules are generally released by one cell type

5 and act to influence proliferation of other cell types (James, R. and Bradshaw, R. A., 1984, *Ann. Rev. Biochem.* 53, 259-292 [see reference I-1 in Experimental Section I, below]). Interest in growth factors has been

heightened by evidence of their potential involvement in neoplasia (Sporn, M. B. and Todaro, G. J., 1980, *N. Eng. J. Med.* 303, 878-880 [reference II-2 in Experimental Section II, below]). The v-sis transforming gene of simian sarcoma virus encodes

a protein that is homologous to the B chain of platelet-derived growth factor (James, R. and Bradshaw, R. A., 1984, *Ann. Rev. Biochem.* 53, 259-292 [I-1], [I-2]). Moreover, a number of oncogenes are homologues of

15 genes encoding growth factor receptors (James, R. and Bradshaw, R. A., 1984, *Ann. Rev. Biochem.* 53, 259-292 [II-1]). Thus, increased understanding of growth factors and their receptor-mediated signal transduction pathways is likely to provide insights into mechanisms of both normal and malignant cell growth.

20 One known family of growth factors affecting connective tissue cells includes acidic fibroblast growth factor (aFGF), basic fibroblast growth factor (bFGF), and the related products of the *hst*^[FGF-5] and *int-2* oncogenes.

Further, it is known that some growth factors, including the following, have heparin-

This invention contemplates the application of methods of protein isolation and recombinant DNA technologies to fulfill such needs and to develop means for producing protein factors of mesenchymal origin, which appear to be related to epithelial cell proliferation processes and which could not be produced otherwise. This invention also contemplates the application of the molecular mechanisms of these factors related to epithelial cell growth processes.

SUMMARY OF THE INVENTION

The present invention relates to developments of protein isolation and recombinant DNA technologies, which include production of novel growth factor proteins affecting epithelial cells, free of other peptide factors. Novel DNA segments and bioassay methods are also included.

The present invention in particular relates to a novel protein having structural and/or functional characteristics of a known family of growth factors which includes acidic fibroblast growth factor (aFGF), basic fibroblast growth factor (bFGF) and the related products of the *hst-1* and *int-2* oncogenes. This new member of the FGF polypeptide family retains the heparin-binding properties of the FGFs but has evolved a unique target cell specificity. This growth factor appears to be specific for epithelial cells and is particularly active on keratinocytes. Therefore, this novel factor has been designated "keratinocyte growth factor" (KGF). Notwithstanding its lack of activity on fibroblasts, since it is the sixth known member of the FGF polypeptide family, KGF may also be referred to as FGF-6.

Accordingly, this invention relates, in part, to purified KGF or KGF-like proteins and

methods for preparing these proteins. Such purified factors may be made by cultivation of human cells which naturally secrete these proteins and application of isolation methods according to the practice of this invention. These proteins can be used for biochemical and biological studies leading, for example, to isolation of DNA segments encoding KGF or KGF-like polypeptides.

10 The present invention also relates to such DNA segments which encode KGF or KGF-like proteins. In a principal embodiment, the present invention relates to DNA segments, which encode KGF-related products, consisting of: human cDNA clones 32 or 49, derived from polyadenylated RNA extracted from the human embryonic lung fibroblast cell line M426; recombinants and mutants of these clones; and related DNA segments which can be detected by hybridization to any of the above human DNA segments, which related segments encode KGF-like proteins or portions thereof.

20 In the practice of one embodiment of this invention, the DNA segments of the invention are capable of being expressed in suitable host cells, thereby producing KGF or KGF-like proteins. The invention also relates to mRNAs produced as the result of transcription of the

sense strands of the DNA segments of this invention.

5 In another embodiment, the invention relates to a recombinant DNA molecule comprising a vector and a DNA of the present invention. These recombinant molecules are exemplified by molecules comprising a KGF cDNA and any of the following vector DNAs: a bacteriophage λ cloning vector (exemplified by λ pCEV9); a DNA sequencing
10 plasmid vector (e.g., a pUC variant); a bacterial gene expression vector (e.g., pKK233-2); or a mammalian gene expression vector (such as pMMT).

In still another embodiment, the invention comprises a cell, preferably a
15 mammalian cell, transformed with a DNA of the invention. Further, the invention comprises cells, including insect cells, yeast cells and bacterial cells such as those of *Escherichia coli* and *B. subtilis*, transformed with DNAs of the invention. According to another embodiment of this aspect of
20 the invention, the transforming DNA is capable of being expressed in the cell, thereby increasing in the cell the amount of KGF or KGF-like protein encoded by this DNA.

25 The primary KGF translation product predicted from its cDNA sequence contains an N-terminal hydrophobic region which likely serves

as a signal sequence for secretion and which is not present in the mature KGF molecule. In a most preferred embodiment of the gene expression aspect of the invention, the cell transformed by the DNA of the invention secretes the protein encoded by that DNA in the (truncated) form that is secreted by human embryonic lung fibroblast cells.

Still further, this invention contemplates KGF or KGF-like proteins produced by expression of a DNA of the invention, or by translation of an RNA of the invention. Preferably, these proteins will be of the secreted form (i.e., lacking an apparent signal sequence). These protein factors can be used for functional studies, and can be purified for additional structural and functional analyses, such as qualitative and quantitative receptor binding assays.

Moreover, the ability to produce large quantities of this novel growth factor by recombinant techniques will allow testing of its clinical applicability in situations where specific stimulation of growth of epithelial cells is of particular importance. Accordingly, this invention includes pharmaceutical compositions comprising KGF or KGF-like polypeptides for use in the treatment of such

conditions, including, for example, healing of wounds due to burns or stimulation of transplanted corneal tissue.

5 According to this embodiment of the invention, the novel KGF-like proteins will be protein products of "unmodified" DNAs and mRNAs of the invention, or will be modified or genetically engineered protein products. As a result of engineered mutations in the DNA
10 sequences, modified KGF-like proteins will have one or more differences in amino acid sequence from the corresponding naturally occurring "wild-type" proteins. According to one embodiment of this aspect of this invention, the modified KGF-
15 like proteins will include "chimeric" molecules comprising segments of amino acid sequences of KGF and at least one other member of the FGF peptide family.

20 Ultimately, given results of analogous successful approaches with other peptide factors having similar properties, development of such chimeric KGF-like polypeptides should lead to superior, "second generation" forms of KGF-like peptides for clinical purposes. These modified
25 KGF-like products might be smaller, more stable, more potent, and/or easier or less expensive to produce, for example.

This invention further comprises novel bioassay methods for determining expression in human cells of the mRNAs and proteins produced from the genes related to DNA segments of the invention. According to one such embodiment, DNAs of this invention may be used as probes to determine steady state levels or kinetics of induction of related mRNAs. The availability of the KGF-related cDNA clones makes it possible to determine whether abnormal expression of this growth factor is involved in clinical conditions characterized by excessive epithelial cell growth, including dysplasia and neoplasia (e.g., psoriasis, or malignant or benign epithelial tumors).

This invention also contemplates novel antibodies made against a peptide encoded by a DNA segment of the invention. In this embodiment of the invention, the antibodies are monoclonal or polyclonal in origin, and are generated using KGF-related polypeptides from natural, recombinant or synthetic chemistry sources.

The antibodies of this invention bind specifically to KGF or a KGF-like protein which includes the sequence of such peptide, preferably when that protein is in its native (biologically active) conformation. These antibodies can be used for detection or purification of the KGF or

[antibodies can be used for detection or
purification of the] KGF or KGF-like protein
factors. In a most preferred embodiment of this
aspect of the invention, the antibodies will
5 neutralize the growth promoting activity of KGF,
thereby enabling mechanistic studies and,
ultimately, therapy for clinical conditions
involving excessive levels of KGF.

BRIEF DESCRIPTION OF THE DRAWINGS

Fig. [I-1] depicts results of heparin-Sepharose affinity chromatography of conditioned medium from M426 human embryonic fibroblasts showing that greater than 90% of the mitogenic activity for mouse keratinocytes (BALB/MK) eluted with 0.6 M NaCl.

Heparin-Sepharose affinity chromatography of conditioned medium from M426 human embryonic fibroblasts. Approximately 150 ml of ultrafiltration retentate derived from five liters of M426 conditioned medium were loaded onto a heparin-Sepharose column (6 ml bed volume) in 1 hr. After washing the column with 150 ml of the equilibration buffer, 20 mM Tris-HCl, pH 7.50/0.3M NaCl, the retained proteins (15% of the total protein in the retentate) was eluted with a modified linear gradient of increasing NaCl concentration. Fraction size was 3.8 ml and flow rate during gradient elution was 108 ml/hr. Two 41 of the indicated fractions were transferred to microtiter wells containing a final volume of 0.2 for assay of ³H-thymidine incorporation in BALB/MK cells as described in the Methods.

Fig. [I-2] illustrates results of further purification of the mitogen from human fibroblasts using HPLC with [and] adsorptive matrix. Panel (A) shows the profile on reversed-phase (C₄) HPLC of BALB/MK mitogenic activity. Panel (B) presents electrophoretic (NaDodSO₄/PAGE) analysis of selected fractions from the C₄ chromatography shown in panel A, demonstrating that the peak HPLC fractions contained a single band on the silver stained gel. Panel (C) is a bar graph of DNA synthesis in BALB/MK cells triggered by the fractions analyzed in Panel B, showing that the relative mitogenic activity correlated well with the intensity of the protein band across the activity profile.]

A, 25 and 2C

Reversed-phase C₄HPLC of BALB/MK mitogenic activity. Active fractions eluted from heparin-Sepharose with 0.6 M NaCl were processed with the Centricon-10 and loaded directly onto a C₄ Vydac column (4.6 x 250mm) which had been equilibrated in 0.1% trifluoroacetic acid/20% acetonitrile (ACN). After washing the column with 4 ml of equilibration buffer, the sample was eluted with a modified linear gradient of increasing % ACN. Fraction size was 0.2 ml and flow rate was 0.5 ml/min. Aliquots for the assay of ³H-thymidine incorporation in BALB/MK cells were promptly diluted 10-fold with 50 µg/ml bovine serum albumin/20 mM Tris-HCl, pH 7.5, and tested at a final dilution of 200-fold. (B) NaDodSO₄/PAGE analysis of selected fractions from the C₄ chromatography shown in panel A. Half of each fraction was dried, redissolved in NaDodSO₄/2-mercaptoethanol, heat denatured and electrophoresed in a 14% polyacrylamide gel which was subsequently stained with silver. The position of each molecular weight marker (mass in kDa) is indicated by an arrow. (C) DNA synthesis in BALB/MK cells triggered by the fractions analyzed in Panel B. Activity is expressed as the fold stimulation over background which was 100 cpm.

Fig. [I-3] presents an alternative purification step to RP-HPLC, using [sieving] chromatography with a (TSK G3000SW GlasPac) column run in aqueous solution near physiological pH. Activity is expressed as the fold stimulation over background which was 100 cpm.

[pH, which resulted in a major peak of mitogenic activity in the BALB/MK bioassay].

Comparison of BALB/MK DNA synthesis in response to TSK-purified mitogen and other growth factors. Incorporation of ^3H -thymidine into trichloroacetic acid-insoluble DNA, expressed as fold stimulation over background, was measured as a function of the concentration of the indicated growth factors.

Background values with no sample added were 150 cpm. Fig. [I-4] illustrates a comparison of BALB/MK DNA synthesis in response to TSK-purified mitogen and other growth factors.

The results represent mean values of two independent experiments. Replicates in each experiment were within 10% of mean values. TSK-purified mitogen, Δ ; EGF, \square ; aFGF, \square ; bFGF, \circ .

Fig. [I-5] shows comparative growth of BALB/MK cells in a chemically defined medium in response to different combinations of growth factors. Cultures were plated at a density of 2.5×10^4 cells per dish on 35 mm Petri dishes precoated with poly-D-lysine/fibronectin in a 1:1 mixture of Eagle's minimal essential medium and Ham's F12 medium supplemented with transferrin, N_2 , Se , O_3 , ethanolamine and the growth factors indicated below. After 10 days, the plates were fixed and stained with Giemsa. Key: a) no growth factor; b) EGF alone; c) insulin alone; d) KGF alone; e) EGF and dialyzed fetal calf serum (final concentration, 10%); f) KGF and EGF; g) KGF and insulin; h) EGF and insulin. Final concentrations of the growth factors were as follows: EGF, 20 ng/ml; insulin, 10 $\mu\text{g}/\text{ml}$; and KGF, 40 ng/ml.

Table I-1 summarizes the results from various purification steps, documenting that sieving chromatography provided a far better recovery of activity than the adsorptive RP-HPLC approach.

Table I-2 recapitulates data on the target cell specificities of various growth factors, demonstrating that the newly isolated factor exhibited a strong mitogenic effect on keratinocytes (BALB/MK) and, in striking contrast, had no detectable effects on fibroblasts or human saphenous vein endothelial cells.

Fig. II-1 presents the nucleotide sequence and deduced amino acid sequence of KGF cDNA, as well as identification of RNAs transcribed from the KGF gene. Panel (A) outlines a schematic representation of human KGF potential asparagine-linked glycosylation site is underlined. The variant polyadenylation signals, AATTAA and AATACA close to the 3' end of the RNA, are boxed.

Fig. 6 outlines a schematic representation of human KGF cDNA clones. Overlapping pCEV9 clones 3a and 49, used in sequence determination, are shown above a diagram of the complete structure in which untranslated regions are depicted by a line and the coding sequence is boxed. The hatched region denotes sequences of the signal peptide. Selected restriction sites are indicated.

Fig. 7 documents the KGF cDNA nucleotide and predicted amino acid sequences. Nucleotides are numbered on the left; amino acids are numbered throughout. The N-terminal peptide sequence derived from purified KGF is underlined. The hydrophobic N-terminal domain is italicized. The potential asparagine-linked glycosylation site is underlined. The variant polyadenylation signals, AATTAA and AATACA close to the 3' end of the RNA, are boxed.

Fig. 8 shows identification of KGF mRNAs by Northern blot analysis. Lanes (a)

Molecular sieving HPLC (TSK 3000SW) chromatography of the BALB/MK mitogenic activity. Approximately 50 μM of a Centricon-processed, 0.6 M NaCl pool from HSAC were loaded onto a LKB GlasPac TSK G 3000SW column (6 x 300 mm), previously equilibrated in 20 mM Tris-HCl, pH 6.8/0.5 M NaCl, and eluted as 0.2 ml fractions at a flow rate of 0.4 ml/min. Aliquots of 2 μM were transferred to microtiter wells containing a final volume of 0.2 ml for assay of ^3H -thymidine incorporation in BALB/MK cells. The elution positions of molecular weight markers (mass in kDa) were as indicated by the arrows.

Fig. 6 outlines a schematic representation of human KGF cDNA clones. Overlapping pCEV9 clones 3a and 49, used in sequence determination, are shown above a diagram of the complete structure in which untranslated regions are depicted by a line and the coding sequence is boxed. The hatched region denotes sequences of the signal peptide. Selected restriction sites are indicated.

Fig. 7 documents the KGF cDNA nucleotide and predicted amino acid sequences. Nucleotides are numbered on the left; amino acids are numbered throughout. The N-terminal peptide sequence derived from purified KGF is underlined. The hydrophobic N-terminal domain is italicized. The potential asparagine-linked glycosylation site is underlined. The variant polyadenylation signals, AATTAA and AATACA close to the 3' end of the RNA, are boxed.

Fig. 8 shows identification of KGF mRNAs by Northern blot analysis. Lanes (a)

and c, poly (A)- selected
M 426 RNA; lanes band d, total
cellular M 426 RNA. Filters were
hybridized with a ³²P-labeled
695 bp Bam HI/Bcl I fragment from
clone 32 (Probe A, Fig. 6), lanes
a and b, or a 541 bp Apa I/Eco RI
fragment from clone 49 (Probe B,
Fig. 6), lanes c and
d.

[nucleotide and predicted amino acid sequences.
(C) Identification of RNA transcripts of KGF
genes by Northern blot analysis.]

Fig. [II-2]⁹ illustrates the topological

5 comparison of the FGF family of related
molecules, including KGF, with emphasis on the
two protein domains that share high homology, the
(shaded boxes)
(hatched boxes)
putative signal peptide sequences, and the two
conserved cysteine residues. (positions labeled with a "c")

10 [Fig. II-3 shows (Northern blot) analyses of expression of KGF-related mRNA in selected
normal human cell lines and tissues, revealing
that a single 2.4 kb transcript was present in
RNA from human embryonic lung fibroblasts and
15 from adult skin fibroblasts, while no transcript
was detected in the (B5/589) epithelial or (HA83)
glial cell lines, or in primary cultures of human
saphenous vein endothelial cells.] Fig. 10 shows
northern blot
analysis of KGF mRNA
in normal human
cell lines and tissues,
and comparison with
mRNA expression of
other growth factors with
known activity on
epithelial cells. Total
cellular RNAs were
isolated by cesium
gradient centrifug-
ation. 10 µg of RNA
were denatured and
electrophoresed in 1%
formaldehyde gels. Follow-
ing milk alkali
denaturation (50 mM
NaOH for 30'), RNA
was transferred to
nitrocellulose filters
using 1M ammonium
acetate as a coelectant.
Filters were hybridized
to a ³²P-labeled cDNA
probe containing the
647 bp Eco RI fragment
from the 5' end of the
KGF coding sequence
(A) or similar probes from
other growth factor DNAs.

20 [Table II-1 summarizes a comparison of
the effect of heparin on KGF mitogenic activity
with effects on other growth factors, showing
that thymidine incorporation into DNA by BALB/MK
cells in response to KGF was inhibited by
heparin, in contrast, to the activities of both
25 aFGF and bFGF which were increased by the same
treatment.]

The following human cell types were
used: squamous cell carcinomas (A 253,
A 388 and A 431); mammary epithelial cells
(B5/589); immortalized bronchial epithelial
15 cells (56 and R1); keratinocytes immortalized
with Ad 12-SV 40; primary human
keratinocytes; neonatal foreskin fibroblasts
(AG 1523); adult skin fibroblasts (50T); and
embryonic lung fibroblasts (WI-38 and M426), and tissues revealing that a
single 2.4 kb transcript was present in RNA from human embryonic
lung fibroblasts and from adult skin fibroblasts, while no transcript was detected in
the (B5/589) epithelial or (HA83) glial cell lines or in primary cultures of human saphenous vein endothelial cells.

DESCRIPTION OF SPECIFIC EMBODIMENTS

This invention relates, in part, to purified KGF or KGF-like proteins and methods for preparing these proteins. A principal embodiment of this aspect of this invention relates to homogeneous KGF characterized by an apparent molecular weight of about 28 kDa based on migration in NaDodSO₄/PAGE, movement as a single peak on reversed-phase high performance liquid chromatography, and a specific activity of at least about 3.4×10^6 units per milligram, and preferably at least about 3.2×10^5 units per milligram, where one unit of activity is defined as that amount which causes half of the maximal possible stimulation of DNA synthesis in certain epithelial (keratinocyte) cells under standard assay conditions outlined below.

To identify novel growth factors specific for epithelial cell types, a clonal

BALB/c mouse keratinocyte cell line, designated ^{Weissman, B.E. and Aaronson, S.A., 1983 Cell 32, 599-606} BALB/MK (I-6) was employed as an indicator cell

to detect such factors. These cells are

dependent for their growth upon an exogenous

source of an epithelial cell mitogen even in ^{Weissman, B.E. and Aaronson, S.A., 1983, Cell 32, 599-606} medium containing serum (I-6). The development

of chemically defined medium for these cells has made it possible to demonstrate that two major

mitogenic pathways are required for BALB/MK proliferation. One involves insulin-like growth factor I (or insulin at high concentration) and the other is satisfied by epidermal growth factor (EGF), transforming growth factor α (TGF α),

acidic fibroblast growth factor (aFGF) or basic fibroblast growth factor (bFGF) (I-7). ^{Falco, J.P., et al., 1988, Oncogene 2, 573-57}

By using BALB/MK as the prototypical epithelial cell line and NIH/3T3 as its fibroblast counterpart, conditioned media from various human cell lines were assayed for new epithelial cell-specific mitogens. These bioassays of this invention enabled the purification to homogeneity of one such novel growth factor, released by a human embryonic lung fibroblast line, and designated herein as keratinocyte growth factor (KGF).

In brief, the bioassay for KGF-like activity under standard conditions comprises the following steps:

(i) Mouse keratinocytes (BALB/MK cells) are grown in culture to confluency and then maintained for 24-72 hr in serum-free medium;

(ii) Following addition of test samples, stimulation of DNA synthesis is determined by incorporation of ³H-thymidine into acid-precipitable DNA.

To determine the [cell target]^{target cell} specificity of a mitogenic growth factor, the DNA synthesis stimulation, expressed as ratio of stimulated synthesis over background incorporation of thymidine in the absence of added test sample, can be compared to analogous stimulation observed in cells other than keratinocytes under the same assay conditions. In such comparisons, KGF mitogenic activity will exhibit marked specificity for the keratinocytes as opposed to fibroblasts (at least about 500-fold greater stimulation) and lesser but significant (at least about 50-fold) greater activity on keratinocytes than on other exemplary epithelial cell types (see Table [I-2] for further data, and Materials and Methods in Experimental Section I for details of the standard conditions of the bioassay).

By employing a method of KGF production involving culturing cells and isolating mitogenic activity, which method comprises ultrafiltration, heparin-Sepharose affinity chromatography (HSAC) and adsorptive reversed-phase high performance liquid chromatography (RP-HPLC) or, alternatively, molecular sieving HPLC (TSK-HPLC), according to the present invention, a quantity was isolated sufficient to permit detailed characterization of the physical and biological properties of this molecule.

To summarize, the method for production of KGF from producing cells such as M426 human embryonic fibroblasts (I-8)^{Aaronson, S.A., and Todaro, G.L. 1968, Virology 36, 254-261}, for example, comprises the following steps:

5 (i) Preparation of conditioned media (e.g., 10 liters) using monolayer cultures cycled from serum-containing to serum-free medium and storing the serum-free harvest at -70°C until further use;

10 (ii) Concentration by ultrafiltration using membranes having a 10 kDa molecular weight cutoff in several successive steps with intervening dilution in buffer (to facilitate removal of low molecular weight materials), followed by optional
15 storage at -70°C;

 (iii) Affinity chromatography on heparin attached to a polymeric support (e.g., Sepharose) with elution by a gradient of increasing NaCl concentration;

20 (iv) Concentration by a factor of at least ten- to twenty-fold with small scale ultrafiltration devices with a 10 kDa molecular weight cutoff (e.g., a Centricon-10 microconcentrator from Amicon) and storage at
25 -70°C.

The next step of the purification process comprises either step (v) or, alternatively, step (vi), as follows:

(v) Reversed-phase HPLC of active fractions (0.6 M NaCl pool) from the previous HSAC step in organic solvent systems;

or,

- 5 (vi) Molecular sieve HPLC (e.g, on a TSK-G3000SW Glas-Pac Column from LKB) in aqueous buffer at near physiological pH (e.g., Tris-HCl, pH 6.8/0.5M NaCl) followed by storage at -70°C.

10 A preparation made by the TSK step (vi) was almost as pure as one obtained from RP-HPLC, as judged by silver-stained NaDodSO₄/PAGE (data not shown); but the TSK approach provided a far better recovery of activity (Table [I-1]).

15 Further, the TSK-purified material had a higher specific activity than the RP-HPLC material. KGF prepared by the TSK procedure above stimulated DNA synthesis in epithelial cells at sub-nanomolar concentrations, but failed to induce any thymidine incorporation into DNA of

20 fibroblasts or endothelial cells at comparable or higher concentrations (up to 5 nM). The activity was sensitive to acid, heat and solvents used in the RP-HPLC step. (See Experimental Section I for data on sensitivities and further details of the

25 production method.)

Using standard methodology well known in the art, an unambiguous amino acid sequence was determined for positions 2-13 from the amino

terminus of the purified KGF, as follows: Asn-Asp-Met-Thr-Pro-Glu-Gln-Met-Ala-Thr-Asn-Val (see Experimental Section I).

5 The present invention also includes DNA segments encoding KGF and KGF-like polypeptides. The DNAs of this invention are exemplified by . DNAs referred to herein as: human cDNA clones 32 and 49 derived from polyadenylated RNA extracted from the human embryonic lung fibroblast cell
10 line M426; recombinants and mutants of these clones; and related DNA segments which can be detected by hybridization to these DNA segments.

As described in Experimental Section II, to search for cDNA clones corresponding to the
15 known portion of the KGF amino acid sequence, two pools of oligonucleotide probes were generated based upon all possible nucleotide sequences encoding the nine-amino acid sequence, Asn-Asp-Met-Thr-Pro-Glu-Gln-Met-Ala. A cDNA
20 library was constructed in a cDNA cloning vector, λ pCEV9, using polyadenylated RNA extracted from the human embryonic lung fibroblast cell line M426 which was the initial source of the growth factor. Screening of the library (9×10^5
25 plaques) with the ^{32}P -labelled oligonucleotides identified 88 plaques which hybridized to both probes.

Of 10 plaque-purified clones that were analyzed, one, designated clone 49, had a cDNA insert of 3.5 kb, while the rest had inserts ranging from 1.8 kb to 2.1 kb. Analysis of the smaller clones revealed several common restriction sites, and sequencing of a representative smaller clone, designated clone 32, along with clone 49, demonstrated that they were overlapping cDNAs (Fig^b [II-1A]). Alignment of the two cDNAs established a continuous sequence of 3.85 kb containing the complete KGF coding sequence. The sense strand DNA nucleotide sequence, and the predicted primary protein sequence encoded, are shown for the full-length composite KGF cDNA sequence in Fig. [II-1B].

These DNAs, cDNA clones 32 and 49, as well as recombinant forms of these segments comprising the complete KGF coding sequence, are most preferred DNAs of this invention.

From the cDNA sequence, it is apparent that the primary KGF₁^[FGF-5] and *hst* translation products contain hydrophobic N-terminal regions which likely serve as signal sequences, based on similarity to such sequences in a variety of other proteins. Accordingly, this N-terminal domain is not present in the purified mature KGF molecule which is secreted by human embryonic fibroblasts.

Furthermore, KGF shares with all other members of the FGF family two major regions of homology, spanning amino acids 65-156 and 162-189 in the predicted KGF sequence, which are
5 separated by short, nonhomologous series of amino acids of various lengths in the different family members. The sequence of the purified form of KGF contains five cysteine residues, two of which are conserved throughout the family of FGF
10 related proteins. Five pairs of basic residues occur throughout the KGF sequence. This same pattern has been observed in other FGF family members.

It should be obvious to one skilled in
15 the art that, by using the DNAs and RNAs of this invention in hybridization methods (such as Southern blot analyses of genomic human DNAs), especially the most preferred DNAs listed herein above, without undue experimentation, it is
20 possible to screen genomic or cDNA libraries to find other KGF-like DNAs which fall within the scope of this invention. Furthermore, by so using DNAs of this invention, genetic markers associated with the KGF gene, such as restriction
25 fragment length polymorphisms (RFLPs), may be identified and associated with inherited clinical conditions involving this or other nearby genes.

This invention also includes modified forms of KGF DNAs. According to a chief embodiment of this aspect of the invention, such modified DNAs encode KGF-like proteins comprising segments of amino acid sequences of KGF and at least one other member of the FGF peptide family. Thus, for example, since there is no significant N-terminal homology between the secreted form of KGF and analogous positions in other FGF-related proteins, polypeptides with novel structural and functional properties may be created by grafting DNA segments encoding the distinct N-terminal segments of another polypeptide in the FGF family onto a KGF DNA segment in place of its usual $[\text{NH}_2]^N$ terminal sequence.

The polypeptide chimeras produced by such modified DNAs are useful for determining whether the KGF NH_2 -terminal domain is sufficient to account for its unique target cell specificity. Studies on chimeras should also provide insights into which domains contribute the different effects of heparin on their biologic activities.

Indeed, the utility of this approach has already been confirmed by the successful engineering and expression of a chimeric molecule in which [about 40] ^{about 40} amino acids from the NH_2 -terminus of the secreted form of KGF (beginning

with the amino terminal cys residue of the mature KGF form, numbered 32 in Fig [II-1],⁷ and ending at KGF residue 78, arg) is linked to [about 140]^{[11/2] about 140} amino acids of the [CO₂]^C-terminal core of aFGF (beginning at residue 39, arg, and continuing to the C-terminal end of the aFGF coding sequence. This chimeric product has a target cell preference for keratinocytes, like KGF, but [lacks susceptibility to] ^[its activity is enhanced by] heparin, a characteristic which parallels that of aFGF rather than KGF. This novel KGF-like growth factor may have advantages in clinical applications where administration of an epithelial-specific growth factor is desirable in the presence of heparin, a commonly used anticoagulant. Further details of the construction of this chimeric molecule and the properties of the polypeptide are described in Experimental Section II.

Other DNAs of this invention include the following recombinant DNA molecules comprising a KGF cDNA and any of the following exemplary vector DNAs: a bacteriophage λ cloning vector (λ pCEV9); a DNA sequencing plasmid vector (a pUC variant); a bacterial expression vector (pKK233-2); or a mammalian expression vector (pMMT/neo). Such recombinant DNAs are exemplified by

constructs described in detail in the
Experimental Sections.

Most preferred recombinant molecules
include the following: molecules comprising the
5 coding sequence for the secreted form of KGF and
a bacterial expression vector (e.g., pKK233-2) or
a cDNA encoding the entire primary translation
product (including the ^N[NH₂]-terminal signal
peptide) and a mammalian expression vector
10 (exemplified by pMMT) capable of expressing
inserted DNAs in mammalian (e.g., NIH/3T3) cells.

Construction of recombinant DNAs
containing KGF DNA and a bacterial expression
vector is described in Experimental Section II.

15 In brief, KGF cDNA was expressed to produce
polypeptide in *E. coli* by placing its coding
sequence under control of the hybrid trk promoter
in the plasmid expression vector pKK233-2 (II-
31). Amman, E. and Brosius, J., 1985, *Gene* 40, 183

20 Construction of recombinant DNAs
comprising KGF DNA and a mammalian vector capable
of expressing inserted DNAs in cultured human or
animal cells, can be carried out by standard gene
expression technology using methods well known in
25 the art for expression of such a relatively
simple polypeptide. One specific embodiment of a
recombinant DNA of this aspect of the present

invention, involving the mammalian vector pMMT, is described further below in this section under recombinant cells of this invention.

DNAs and sense strand RNAs of this invention can be employed, in conjunction with protein production methods of this invention, to make large quantities of substantially pure KGF or KGF-like proteins. Substantially pure KGF protein thus produced can be employed, using well-known techniques, in diagnostic assays to determine the presence of receptors for this protein in various body fluids and tissue samples.

Accordingly, this invention also comprises a cell, preferably a bacterial or mammalian cell, transformed with a DNA of the invention, wherein the transforming DNA is capable of being expressed. In a preferred embodiment of this aspect of the invention, the cell transformed by the DNA of the invention produces KGF protein in a fully mitogenic form. Most preferably, these proteins will be of a secreted form (i.e., lacking an apparent signal sequence). These protein factors can be used for functional studies, and can be purified for additional biochemical and functional analyses, such as qualitative and quantitative receptor binding assays.

Recombinant *E. coli* cells have been constructed in a bacterial expression vector, pKK233-2, for production of KGF, as detailed in Experimental Section II. In summary, several
5 recombinant bacterial clones were tested for protein production by the usual small scale methods. All recombinants tested synthesized a protein that was recognized by antibodies raised against an amino-terminal KGF peptide (see
10 below). One recombinant was grown up in a one liter culture which produced recombinant KGF that efficiently stimulated thymidine incorporation into DNA of BALB/MK keratinocyte cells, but was only marginally active on NIH/3T3 fibroblasts.
15 Half-maximal stimulation of the BALB/MK cells in the standard keratinocyte bioassay was achieved with a concentration of between 2 to 5 ng/ml, compared to a concentration of 10 to 15 ng/ml for KGF purified from M426 cells.

20 One liter of bacterial cells yielded approximately 50 μ g of Mono-S purified recombinant KGF. It will be apparent to those skilled in the art of gene expression that this initial yield can be improved substantially
25 without undue experimentation by application of a variety known recombinant DNA technologies.

Recombinant mammalian (NIH/3T3 mouse) cells have also been constructed using the entire

KGF cDNA coding sequence (including the NH₂-terminal signal peptide) and the vector pMMT/neo, which carries mouse metallothionine (MMT) promoter and the selective marker gene for neomycin resistance. The cells are being evaluated for KGF production, particularly for secretion of the mature form (lacking signal peptide) produced by human fibroblasts, using bioassays of the present invention. This same vector and host cell combination has been used successfully to express several other similar recombinant polypeptides, including high levels of Platelet-Derived Growth Factor (PDGF) A and B chains (II-32).^{Sakai, R.K., Scharf, S., Faloona, F., Mullis, K.B., Norn, G.T., Erlich, H.A. and Arnheim, N. (1985) Science 230, 1358-135X} Accordingly, it will be recognized by those skilled in the art that high yields of recombinant KGF can be achieved in this manner, using the aforementioned recombinant DNAs and transformed cells of this invention.

Ultimately, large-scale production can be used to enable clinical testing in conditions requiring specific stimulation of epithelial cell growth. Materials and methods for preparing pharmaceutical compositions for administration of polypeptides topically (to skin or to the cornea of the eye, for example) or systemically are well known in the art and can be adapted readily for

administration of KGF and KGF-like peptides without undue experimentation.

This invention also comprises novel antibodies made against a peptide encoded by a DNA segment of the invention. This embodiment of the invention is exemplified by several kinds of antibodies which recognize KGF. These have been prepared using standard methodologies well known in the art of experimental immunology, as outlined in Experimental Section II. These antibodies include: monoclonal antibodies raised in mice against intact, purified protein from human fibroblasts; polyclonal antibodies raised in rabbits against synthetic peptides with sequences based on amino acid sequences predicted from the KGF cDNA sequence [exemplified by a peptide with the sequence of KGF residues 32-45]⁴⁵ plus an R at its C terminus, namely, NDMTPEQMATNVR (using standard one-letter code for amino acid sequences; see Fig. [II-1]); polyclonal antibodies raised in rabbits against both naturally secreted KGF from human fibroblasts and recombinant KGF produced in *E. coli* (see above).

All tested antibodies recognize the recombinant as well as the naturally occurring KGF, either in a solid-phase (ELISA) assay and/or in a Western blot. Some exemplary antibodies, which are preferred antibodies of this invention,

appear to neutralize mitogenic activity of KGF in the BALB/MK bioassay.

Fragments of antibodies of this invention, such as Fab or F(ab)' fragments, which retain antigen binding activity and can be prepared by methods well known in the art, also fall within the scope of the present invention. Further, this invention comprises pharmaceutical compositions of the antibodies of this invention, or active fragments thereof, which can be prepared using materials and methods for preparing pharmaceutical compositions for administration of polypeptides that are well known in the art and can be adapted readily for administration of KGF and KGF-like peptides without undue experimentation.

These antibodies, and active fragments thereof, can be used, for example, for detection of KGF in bioassays or for purification of the protein factors. They may also be used in approaches well known in the art, for isolation of the receptor for KGF, which, as described in Experimental Section II, appears to be distinct from those of all other known growth factors.

Those preferred antibodies, and fragments and pharmaceutical compositions thereof, which neutralize mitogenic activity of KGF for epithelial cells, as indicated by the

BALB/MK assay, for instance, may be used in the treatment of clinical conditions characterized by excessive epithelial cell growth, including dysplasia and neoplasia (e.g., psoriasis, or malignant or benign epithelial tumors).

This invention further comprises novel bioassay methods for detecting the expression of genes related to DNAs of the invention. In some exemplary embodiments, DNAs of this invention were used as probes to determine steady state levels of related mRNAs. Methods for these bioassays of the invention, using KGF DNAs, and standard Northern blotting techniques, are described in detail in Experimental Section II.

One skilled in the art will recognize that, without undue experimentation, such methods may be readily applied to analysis of gene expression for KGF-like proteins, either in isolated cells or various tissues. Such bioassays may be useful, for example, for identification of various classes of tumor cells or genetic defects in the epithelial growth processes.

Without further elaboration, it is believed that one of ordinary skill in the art, using the preceding description, and following the methods of the Experimental Sections below, can utilize the present invention to its fullest

extent. The material disclosed in the
Experimental Sections, unless otherwise
indicated, is disclosed for illustrative purposes
and therefore should not be construed as being
5 limitive in any way of the appended claims.

EXPERIMENTAL SECTION I

IDENTIFICATION AND CHARACTERIZATION OF A NOVEL GROWTH FACTOR SPECIFIC FOR EPITHELIAL CELLS

This section describes experimental work
5 leading to identification of a growth factor
specific for epithelial cells in conditioned
medium of a human embryonic lung fibroblast cell
line. The factor, provisionally termed
keratinocyte growth factor (KGF) because of its
10 predominant activity on this cell type, was
purified to homogeneity by a combination of
ultrafiltration, heparin-Sepharose affinity
chromatography and hydrophobic chromatography on
a C₈ reversed-phase HPLC column, according to
15 methods of this invention. KGF was found to be
both acid and heat labile, and consisted of a
single polypeptide chain with an apparent
molecular weight of approximately 28,000 daltons.
Purified KGF was a potent mitogen for epithelial
20 cells, capable of stimulating DNA synthesis in
quiescent BALB/MK epidermal keratinocytes by more
than 500-fold with activity detectable at 0.1 nM
and maximal at 1.0 nM. Lack of mitogenic
activity on either fibroblasts or endothelial
25 cells indicated that KGF possessed a target cell
specificity distinct from any previously
characterized growth factor. Microsequencing

revealed an amino-terminal sequence containing no significant homology to any known protein. The release of this novel growth factor by human embryonic fibroblasts indicates that KGF plays a role in mesenchymal stimulation of normal epithelial cell proliferation.

METHODS AND MATERIALS

Preparation of Conditioned Media. An early passage of M426 human embryonic fibroblasts (I-8)^{Aaronson, S.A. and Todaro, G.L. 1968, Virology 36, 254-261} was plated onto 175 cm² T-flasks and grown to confluence over 10-14 days in Dulbecco's modified Eagle's medium (DMEM; GIBCO) supplemented with 10% calf serum (GIBCO). Once confluent, the monolayers were cycled weekly from serum-containing to serum-free medium, the latter consisting of DMEM alone. The cells were washed twice with 5 ml of phosphate buffered saline prior to addition of 20 ml of DMEM. After 72 hrs, culture fluids were collected and replaced with 35 ml of serum-containing medium. The conditioned medium was stored at -70°C until further use.

Ultrafiltration. Approximately ten liters of conditioned medium were thawed, prefiltered through a 0.50 micron filter (Millipore HAWP 142 50) and concentrated to 200

ml using the Pellicon cassette system (Millipore
XX42 00K 60) and a cassette having a 10 kDa
molecular weight cutoff (Millipore PTGC 000 05).
After concentration, the sample was subjected to
5 two successive rounds of dilution with one liter
of 20 mM Tris-HCl, pH 7.5/0.3M NaCl, each
followed by another step of ultrafiltration with
the Pellicon system. Activity recovered in the
retentate was either immediately applied to
10 heparin-Sepharose resin or stored at -70°C.

Heparin-Sepharose Affinity

Chromatography (HSAC). The retentate from
ultrafiltration was loaded onto heparin-Sepharose
resin (Pharmacia) which had been equilibrated in
15 20 mM Tris-HCl, pH 7.5/0.3 M NaCl. The resin was
washed extensively until the optical density had
returned to baseline and then subjected to a
linear-step gradient of increasing NaCl
concentration. After removing aliquots from the
20 fractions for the thymidine incorporation
bioassay, selected fractions were concentrated
ten- to twenty-fold with a Centricon-10
microconcentrator (Amicon) and stored at -70°C.

Reversed-Phase HPLC (RP-HPLC). Active
25 fractions (0.6 M NaCl pool) from the HSAC were
thawed, pooled and further concentrated with the
Centricon-10 to a final volume of $\leq 200 \mu\text{l}$. The
sample was loaded onto a Vydac C₄ HPLC column (The

Separations Group, Hesperia, CA) which had been equilibrated in 0.1% trifluoroacetic acid (TFA, Fluka)/20% acetonitrile (Baker, HPLC grade) and eluted with a linear gradient of increasing acetonitrile. Aliquots for the bioassay were immediately diluted in a 10-fold excess of 50 µg/ml BSA (Fraction V, Sigma)/20 mM Tris-HCl, pH 7.5. The remainder of the sample was dried in a Speed-Vac (Savant) in preparation for structural analysis.

Molecular Sieve HPLC. Approximately 50 µl of the twice concentrated heparin-Sepharose fractions were loaded onto a TSK-G3000SW Glas-Pac Column (LKB) which had been equilibrated in 20 mM Tris-HCl, pH 6.8/0.5M NaCl. The sample was eluted in this buffer at a flow rate of 0.4 ml/min. After removing aliquots for the bioassay, the fractions were stored at -70°C.

NaDodSO₄-Polyacrylamide Gel

Electrophoresis (NaDodSO₄/PAGE). Polyacrylamide gels were prepared with NaDodSO₄ according to the procedure of Laemmli (I-9) ^{Laemmli, U.K., 1970, Nature 227, 680-685}. Samples were boiled for 3 min in the presence of 2.5% 2-mercaptoethanol (vol/vol). The gels were fixed and stained with silver (I-10) ^{Merril, C.R., et al., 1981, Science 211, 1437-1438} using the reagents and protocol from BioRad. Molecular weight markers were from Pharmacia.

DNA Synthesis Stimulation. Ninety-six well microtiter plates (Falcon No. 3596) were precoated with human fibronectin (Collaborative Research) at $1 \mu\text{g}/\text{cm}^2$ prior to seeding with

5 BALB/MK cells. Once confluent, the cells were maintained for 24-72 hr in serum-free medium containing $5 \mu\text{g}/\text{ml}$ transferrin (Collaborative Research) and 30 nM Na_2SeO_3 (Baker).

Incorporation of ^3H -thymidine ($5 \mu\text{Ci}/\text{ml}$ final

10 concentration, NEN) into DNA was measured during a 6 hr period beginning at 16 hrs following addition of samples. The assay was terminated by washing the cells once with ice cold phosphate-buffered saline and twice with 5% trichloroacetic

15 acid. The precipitate was redissolved in 0.25 M NaOH , transferred into liquid scintillation fluid (Biofluor, NEN) and counted.

Stimulation of DNA synthesis was monitored as described above for BALB/MK cells on

20 a variety of other cell lines. NIH/3T3 fibroblasts ([I-11])^{Jainchill, J.L., Aaronsen, S.A. and Todaro, G.J., 1969, J. Virol. 4, 549-553} were available from the

National Institutes of Health, while CCL208

Rhesus monkey bronchial epithelial cells ([I-12])^{Caputo, J.L., Hay, R.J. and Williams, C.D., 1979, In Vitro 15, 222-223} were obtained from the American Type Culture

25 Collection. The B5/589 human mammary epithelial cell line, prepared as described in ([I-13])^{Stampfer, M.R. and Bartley, J.C., 1985, Proc. Natl. Acad. Sci. USA 82, 2394-2398} was obtained from Martha Stampfer (University of

Lawrence Berkeley Laboratory
California, Berkeley). The mammary cells were

grown in RPMI 1640 supplemented with 10% fetal calf serum and 4 ng/ml EGF. When maintained in serum-free conditions, the basal medium was DMEM.

Primary cultures of human saphenous vein

5 endothelial cells were prepared and maintained as described elsewhere ([I-14]⁹). Epidermal growth factor and insulin were from Collaborative

Shoreff, J. B., Fairchild, R. D., Albous, R. A., Cruess, D. F. and Rich, N. M., 1986, J. Surgical Res. 41, 463-472

Research. Acidic FGF and bFGF were obtained from California Biotechnology, Inc. Recombinant TGF α was obtained from Genentech, Inc. Media and serum were either from GIBCO, Biofluids, Inc. or the NIH media unit.

Proliferation Assay. Thirty-five mm culture dishes were precoated sequentially with poly-D-lysine (20 $\mu\text{g}/\text{cm}^2$) (Sigma) and human fibronectin, and then seeded with approximately 2.5 x 10⁴ BALB/MK cells. The basic medium was a 1:1 mixture of Eagle's low Ca²⁺ minimal essential medium and Ham's F-12 medium, supplemented with 5 $\mu\text{g}/\text{ml}$ transferrin, 30 nM Na₂SeO₃ and 0.2 mM ethanolamine (Sigma). Medium was changed every 2 or 3 days. After 10 days, the cells were fixed in formalin (Fisher Scientific Co.) and stained with Giemsa (Fisher Scientific Co.).

25 Protein microsequencing. Approximately 4 μg (-150 pmol) of protein from the active fractions of the C₄ column were redissolved in 50% TFA and loaded onto an Applied Biosystems gas-

phase protein sequenator. Twenty rounds of Edman degradation were carried out and identifications of amino acid derivatives were made with an automated on-line HPLC (Model 120A, Applied Biosystems).

RESULTS

Growth Factor Detection and Isolation.

Preliminary screening of conditioned media from various cell lines indicated that media from some fibroblast lines contained mitogenic activities detectable on both BALB/MK and NIH/3T3 cells.

Whereas boiling destroyed the activity on BALB/MK, mitogenic activity on NIH/3T3 remained intact. Based on the known heat stability of EGF

(I-15) and TGF α (I-16), it was reasoned that the BALB/MK mitogenic activity might be due to an agent different from these known epithelial growth factors.

Cohen, S., 1962, J. Biol. Chem. 237, 1555-1562
DeLarco, J. E. and Todaro, G. L., 1978, Proc. Natl. Acad. Sci. USA 75, 4001-4005

M426, a human embryonic lung fibroblast line, was selected as the most productive source of this activity for purification of the putative growth factor(s). Ultrafiltration with the Pellicon system provided a convenient way of reducing the sample volume to a suitable level for subsequent chromatography. Various combinations of sieving, ion exchange and

isoelectric focusing chromatography were tried during the development of a purification scheme, but all resulted in unacceptably low yields.

On the other hand, heparin-Sepharose affinity

5 chromatography (HSAC), which has been employed in

the purification of other growth factors [I-17] -- Raines, E.W. and Ross, R., 1982, J. Biol. Chem. 257, 5154-5160
Shing, Y., Folkman, J., Sullivan, R., Butterfield, C., Myer, J., and Klagsbrun, M., 1984, Science 223, 1296-1299
[I-22], proved to be useful as an early

purification step in the present invention. Maciag, T., et al., 1984, Science 225, 932-935

While estimates of recovered specific activity Conn, G. and Hatcher, V.B. (1981) Biochem. Biophys. Res. Comm. 124, 262-268
10 were uncertain at this stage because of the Lobb, R. R. and Felt, J. W., 1984, Biochemistry 23, 6295-6299
likely presence of other factors, the apparent
yield of activity was 50-70% with a corresponding
enrichment of approximately 1000 fold.

As shown in Fig. [I-1], greater than 90%
15 of the BALB/MK mitogenic activity eluted from the
HSAC column with 0.6M NaCl. This peak of
activity was not associated with any activity on
NIH/3T3 cells (data not shown). A much smaller
peak of BALB/MK mitogenic activity consistently
20 emerged with 0.8 - 1.2M NaCl.

Due to the reproducibility of the HSAC
pattern, active fractions could be identified
presumptively on the basis of the gradient and
optical density profile. Prompt concentration of
25 10-20 fold with the Centricon-10 was found to be
essential for stability, which could be
maintained subsequently at -70°C for several
months.

Final purification was achieved by RP-HPLC with a C₄ Vydac column, a preparative method suitable for amino acid sequence analysis. While the yield of activity from the C₄ step was usually only a few percent, this loss could be attributed to the solvents employed. In other experiments, exposure to 0.1% TFA/50% acetonitrile for 1 hr at room temperature reduced the mitogenic activity of the preparation by 98%. Nonetheless, as shown in Fig. [I-2]^{2A}, a single peak of BALB/MK stimulatory activity was obtained, coinciding with a distinct peak in the optical density profile. The peak fractions produced a single band upon NaDodSO₄/PAGE and silver staining of the gel (Fig. [I-2B]), and the relative mitogenic activity of each tested fraction (Fig. [I-2C]) correlated well with the intensity of the bands across the activity profile.

An alternative purification step to the HPLC technique described above, using sieving chromatography with a TSK G3000SW GlasPac column run in aqueous solution near physiologic pH, resulted in a major peak of activity in the BALB/MK bioassay (Fig. [I-3]). This preparation was almost as pure as the one obtained from RP-HPLC as judged by silver-stained NaDodSO₄/PAGE (data not shown) but provided a far better recovery of activity (Table [I-1]). The TSK-

purified material was used routinely for biological studies as it had a higher specific activity.

In both types of purified preparations (i.e., purified by HPLC or molecular sieving), the profile of mitogenic activity was associated with a distinct band on NaDodSO₄/PAGE which appeared to be indistinguishable in the two preparations. INSERT Table 1 from page 87

Physical and Biological Characterization of the Growth Factor. The purified factor had an estimated molecular weight of about 28 kDa based on NaDodSO₄/PAGE under reducing (Fig. [I]^{A-2C}₂) and non-reducing conditions (data not shown). This value was in good agreement with its elution position on two different sizing columns run in solvents expected to maintain native conformation (TSK-G3000-SW, Fig. [I]₃, and superose-12, data not shown). From these data, the mitogen appears to consist of a single polypeptide chain with a molecular weight of 25-30 kDa.

The heat and acid lability of the mitogenic activity were demonstrated using the BALB/MK mitogenesis bioassay. While activity was unaffected by a 10 min incubation at 50°C, it was reduced by 68% after 10 min at 60°C and was undetectable after 3 min at 100°C. Exposure to 0.5M acetic acid for 60 min at room temperature

resulted in a decline in activity to 14% of the control. In comparison, the mitogenic activity of the known growth factor, EGF, was not diminished by any of these treatments.

5 The dose response curve for the purified growth factor depicted in Fig. [I-4] illustrates that as little as 0.1 nM led to a detectable stimulation of DNA synthesis. Thus, the activity range was comparable to that of the other growth
10 factors analyzed to date. A linear relationship was observed in the concentration range 0.1 - 1.0 nM with maximal stimulation of 600 fold observed at 1.0 nM. The novel factor consistently induced a higher level of maximal thymidine incorporation
15 than EGF, aFGF, or bFGF in the BALB/MK keratinocytes (Fig. [I-4]).

 The distinctive target cell specificity of this factor was demonstrated by comparing its activities on a variety of cell types with those
20 of other growth factors known to possess epithelial cell mitogenic activity. As shown in Table [I-2], the newly isolated factor exhibited a strong mitogenic effect on BALB/MK but also induced demonstrable incorporation of thymidine
25 into DNA of the other epithelial cells tested. In striking contrast, the factor had no detectable mitogenic effects on mouse (or human,

data not shown) fibroblasts or human saphenous vein endothelial cells.

By comparison, none of the other known growth factors appeared to preferentially stimulate keratinocytes. TGF α and EGF showed potent activity on fibroblasts, while the FGFs were mitogenic for endothelial cells as well as fibroblasts (Table [I-]2). Because of its specificity of epithelial cells and the sensitivity of keratinocytes in particular, the novel mitogen was provisionally designated as keratinocyte growth factor (KGF).

To establish that KGF not only would stimulate DNA synthesis but would also support sustained cell growth, the ability of BALB/MK cells to grow in a fully-defined, serum-free medium supplemented with this growth factor was assessed. As shown in Fig. [I-]5, KGF served as an excellent substitute for EGF but not insulin (or insulin-like growth factor I) in this chemically defined medium. Thus, KGF appears to act through the major signalling pathway shared by EGF, aFGF and bFGF for proliferation of BALB/MK cells.

Insert Table 2, from p 88

Microsequencing Reveals a Unique N-terminal

Amino Acid Sequence of KGF. To further characterize the growth factor, approximately 150 pmol of C₄-purified material were subjected to amino acid sequence analysis. A single sequence

was detected with unambiguous assignments made for cycles 2-13, as follows: X-Asn-Asp-Met-Thr-Pro-Glu-Gln-Met-Ala-Thr-Asn-Val. High background noise precluded an assignment for the first position which is, therefore, indicated by an X.

A computer search using the FASTP ^{Lipman, D.J. and Pearson, R.W., 1985, Science 227, 1435-1441} program ([I-24]) revealed that the N-terminal amino acid sequence of KGF showed no significant homology to any protein in the National Biomedical Research Foundation data bank, thus supporting the novelty of this epithelial growth factor.

DISCUSSION

The studies described in this Experimental Section identified a human growth factor which has a unique specificity for epithelial cells. By employing ultrafiltration, HSAC and RP-HPLC or TSK sieving chromatography according to the present invention, a quantity sufficient to permit detailed characterization of the physical and biological properties of this molecule was isolated.

A single silver-stained band corresponding to a molecular weight of about 28,000 daltons was detected in the active fractions from RP-HPLC, and the intensity of the

band was proportional to the level of mitogenic activity in these fractions. A band indistinguishable from that obtained by RP-HPLC was seen in the active fractions from TSK chromatography. The purified protein stimulated DNA synthesis in epithelial cells at sub-nanomolar concentrations, but failed to induce any thymidine incorporation in fibroblasts or endothelial cells at comparable or higher concentrations (up to 5 nM). This distinctive target cell specificity combined with the single novel N-terminal amino acid sequence determined from the purified molecule lead to the conclusion that KGF represents a new growth factor.

In a chemically defined medium the purified factor was able to complement the insulin-like growth factor I/insulin growth requirement of BALB/MK cells and therefore must act through a signal transduction pathway shared with EGF, TGF α and the FGFs. Moreover, the new factor was more potent than any of the known epithelial cell mitogens in stimulating thymidine incorporation in BALB/MK cells. Preliminary evidence indicates that this factor is also capable of supporting proliferation of secondary cultures of human keratinocytes (data not shown).

Handling and storage of KGF were problematical during its purification. Besides

its inherent lability to acid and heat, it was unstable to lyophilization or dialysis. After HSAC, complete loss of activity occurred within 24 hr despite the use of carrier proteins, heparin, protease inhibitors, siliconized tubes or storage at either 4° or -20°C. Only concentrating the sample at this stage could preserve its activity.

Furthermore, in order to transfer the dried, purified factor it was necessary to utilize either strong acid or detergent, consistent with an adsorptive tendency or insolubility. Thus, for preservation of activity, the purified factor was maintained in solution at high concentrations at -70°C where it remained stable for several months.

The ability of KGF to bind heparin may signify a fundamental property of this factor that has a bearing on its function *in vivo*. Growth

factors with heparin-binding properties include aFGF ([I-20]--[I-22]), bFGF ([I-19], [I-22]), 124, 262-268
 Mauer, T., et al., 1984, *Science* 225, 932-938 Conn, E. and Hatcher, V.B. (1984) *Biochem. Biophys. Res. Com.*
 Lobb, R.R. and Iaff, J.W., 1984, *Biochemistry* 23, 6245-6249 Gaspardowicz, D., et al., 1984, *Proc. Natl. Acad. Sci. USA* 81, 6963-6967
 granulocyte/macrophage colony stimulating factor (Roberts, R., et al., 1988, *Nature* 332, 376-378)
 and interleukin 3. ([I-25]) Each of these is produced by stromal cells ([I-25]--[I-27]). Such factors appear to be deposited in the extracellular matrix, or on proteoglycans coating the stromal cell surface ([I-25], [I-28]). It has been postulated that their storage, release and

contact with specific target cells are regulated

by this interaction (I-25), (I-28). While

mesenchymal-derived effectors of epithelial cell proliferation have also been described

(I-29)-(I-31), their identities have not been elucidated. Its heparin-binding properties, release by human

embryonic fibroblast stromal cells, and

epithelial cell tropism provide KGF with all of

the properties expected of such a paracrine

mediator of normal epithelial cell growth.

The partial amino acid sequence

determined for this new growth factor has enabled

molecular cloning of its coding sequence and

determination of its structural relationship to

known families of growth factors, as described in

Experimental Section II, below.

REFERENCES FOR EXPERIMENTAL SECTION I

- I-1. James, R. and Bradshaw, R.A. (1984) *Ann. Rev. Biochem.* 53, 259-292.
- I-2. Doolittle, R.F., Hunkapiller, M.W., Hood, L.E., Devare, S.G., Robbins, K.C., Aaronson, S.A. and Antoniades, M.N. (1983) *Science* 221, 275-277.
- I-3. Waterfield, M.D., Scrace, G.J., Whittle, N., Strooband, P., Johnson, A., Wasteton, A., Westermarck, B., Heldin,

- [C.-H., Huang, J.S. and Deuel, T.F.
(1983) *Nature* 304, 35-39.
- I-4. Hunter, T. and Cooper, J.A. (1985) *Annu. Rev. Biochem.* 54, 897-930.
- 5 I-5. Wright, N. and Allison, M. (1984) *The Biology of Epithelial Cell Populations* (Oxford University Press, New York) Vol. 1, pp. 3-5.
- I-6. Weissman, B.E. and Aaronson, S.A. (1983)
10 *Cell* 32, 599-606.
- I-7. Falco, J.P., Taylor, W.G., DiFiore, P.P., Weissman, B.E., and Aaronson, S.A. (1988) *Oncogene* 2, 573-578.
- I-8. Aaronson, S.A. and Todaro, G.J. (1968)
15 *Virology* 36, 254-261.
- I-9. Laemmli, U.K. (1970) *Nature* 227, 680-685.
- I-10. Merrill, C.R., Goldman, D., Sedman, S.A. and Ebert, M.H. (1981) *Science* 211, 1437-1438.
- 20 I-11. Jainchill, J.L., Aaronson, S.A. and Todaro, G.J. (1969) *J. Virol.* 4, 549-553.
- I-12. Caputo, J.L., Hay, R.J. and Williams, C.D. (1979) *In Vitro* 15, 222-223.
- I-13. Stampfer, M.R. and Bartley, J.C. (1985)
25 *Proc. Natl. Acad. Sci. USA* 82, 2394-2398.]

- [I-14. Sharefkin, J.B., Fairchild, K.D., Albus, R.A., Cruess, D.F. and Rich, N.M. (1986) *J. Surgical Res.* 41, 463-472.
- I-15. Cohen, S. (1962) *J. Biol. Chem.* 237, 1555-1562.
- 5 I-16. DeLarco, J.E. and Todaro, G.J. (1978) *Proc. Natl. Acad. Sci. USA* 75, 4001-4005.
- I-17. Raines, E.W. and Ross, R. (1982) *J. Biol. Chem.* 257, 5154-5160.
- 10 I-18. Shing, Y., Folkman, J., Sullivan, R., Butterfield, C., Murray, J. and Klagsburn, M. (1984) *Science* 223, 1296-1299.
- I-19. Gospodarowicz, D., Cheng, J., Lui, G.-M., Baird, A. and Bohlen, P. (1984) *Proc. Natl. Acad. Sci. USA* 81, 6963-6967.
- 15 I-20. Maciag, T., Mehlman, T., Friesel, R. and Schreiber, A.B. (1984) *Science* 225, 932-935.
- 20 I-21. Conn, G. and Hatcher, V.B. (1984) *Biochem. Biophys. Res. Comm.* 124, 262-268.
- I-22. Lobb, R.R. and Fett, J.W. (1984) *Biochemistry* 23, 6295-6299.
- I-23. Bradford, M. (1976) *Anal. Biochem.* 72, 248-254.]
- 25

- [I-24. Lipman, D.J. and Pearson, R.W. (1985)
Science 227, 1435-1441.
- I-25. Roberts, R., Gallagher, J., Spooncer,
 5 E., Allen, T.D., Bloomfield, F. and
 Dexter, T.M. (1988) *Nature* 332, 376-378.
- I-26. Libermann, T.A., Friesel, R., Jaye, M.,
 Lyall, R.M., Westermarck, B., Drohen, W.,
 Schmidt, A., Maciag, T. and
 10 Schlessinger, J. (1987) *EMBO J.*, 6, 1627-
 1632.
- I-27. Shipley, G.D., Sternfeld, M.D., Coffey,
 R.J. and Pittelkow, M.R. (1988) *J. Cell*
Biochem. Supp 12A, 125, abstr. C420.
- I-28. Vlodavsky, I., Folkman, J., Sullivan,
 15 R., Fridman, R., Ishai-Michaeli, R.,
 Sasse, J. and Klagsburn, M. (1987) *Proc.*
Natl. Acad. Sci. USA 84, 2292-2296.
- I-29. Gilchrest, B.A., Karassik, R.L.,
 Wilkins, L.M., Vrabel, M.A. and Maciag,
 20 T. (1983) *J. Cell Physiol.* 117, 2325-240.
- I-30. Chan, K.Y. and Haschke, R.H. (1983) *Exp.*
Eye Res. 36, 231-246.
- I-31. Stiles, A.D., Smith, B.T. and Post, M.
 (1986) *Exp. Lung Res.* 11, 165-177.]

EXPERIMENTAL SECTION II

CDNA SEQUENCE OF A NOVEL EPITHELIAL CELL SPECIFIC GROWTH FACTOR DEFINES A NEW MEMBER OF THE FGF FAMILY

5 Work in the previous Experimental
Section I identified and purified a novel
heparin-binding growth factor, designated
keratinocyte growth factor (KGF), which is
particularly active on keratinocytes and appears
10 to be specific for epithelial cells. This second
Experimental Section describes the isolation and
characterization of cDNA clones encoding KGF,
using synthetic oligonucleotides, based upon the
experimentally determined $[\text{NH}_2]^N$ -terminal amino acid
15 sequence, as hybridization probes. Nucleotide
sequence analysis identified a 582-bp open
reading frame which would code for a 194-amino
acid polypeptide that is between 41% and 33%
identical to the heparin-binding acidic and basic
20 fibroblast growth factors (FGFs), and the related
products of the $[\text{FGF-5}]$ *hst-1* and *int-2* oncogenes. The KGF
gene RNA transcript is expressed in normal
fibroblasts of both embryonic and adult origin,
but not in epithelial, endothelial or glial
25 cells. Thus, KGF appears to be normally
expressed by the mesenchyme, indicating a role in
the regulation of epithelial cell proliferation.

MATERIALS AND METHODS

Isolation of cDNA clones. The

purification and N-terminal sequencing of KGF has

been previously described (see Experimental

5 Section I, above and [II-3]). Pools (50 pmole) of deoxyoligonucleotides described under Results

were 5' end-labelled using 83 pmole of γ - ^{32}P -ATP

(3000 Ci/mmole, Amersham) and 10 units of T4

polynucleotide kinase. The recombinant phage

10 carrying cDNA clones were replica plated onto

nitrocellulose filters and hybridized with

^{32}P -labelled deoxyoligonucleotides in 20%

formamide, 10% dextran sulphate, 10 mM Tris-HCl

(pH 7.5), 8 x SSC, 5x Denhardt's and 50 $\mu\text{g/ml}$

15 denatured salmon sperm DNA, overnight at 42°C.

Filters were washed in 0.5 x SSC, 0.1% SDS at

50°C and exposed to Kodak X-omat AR film.

DNA sequencing. The nucleotide sequence

of the KGF cDNA was determined by the dideoxy

20 chain termination method ([II-26]), of overlapping

restriction fragments, subcloned into pUC vectors

([II-27]) Hansch-Perron, C., Vieira, J. and Messing, J., 1985, Gene 33, 103-119

Construction of a bacterial expression

vector for KGF cDNA. KGF cDNA encoding the

25 mature, secreted form of the polypeptide was

placed under control of the hybrid trk promoter

in the plasmid expression vector pKK233-2 (II-31), as follows. To accomplish this, a specific length of KGF cDNA that contained the information to code for the mature KGF molecule (i.e.,

5 without its signal peptide) was amplified using

the polymerase chain reaction (PCR) technique (II-32). The fragment was directionally inserted between two sites in the vector, namely the *Nco*I

site, made blunt ended by *S*1 nuclease digestion,

10 and the *Hind*III site, using standard recombinant DNA methodology. The ends of the KGF cDNA

produced by the PCR method were as follows: the

5' end was blunt and began with an ATG codon,

followed by the codon TGC for cys residue, number

15 33, which is the amino terminal residue of the mature form of KGF (see Fig. [II-1]), and then the entire KGF coding sequence. The stop codon, TAA, and the four bases immediately following, TTGC, were also included on the 3' end of the cDNA.

20 The primer used in the PCR method to direct DNA synthesis to the desired position on the 3' end of the cDNA included a *Hind*III site for insertion of the amplified cDNA into the vector DNA.

Production of antibodies against KGF and

25 KGF-related peptides. Monoclonal antibodies were raised in mice against intact, purified protein from human fibroblasts using 5 or more

subcutaneous injections. Test bleeds were screened with a solid-phase (ELISA) assay using highly purified KGF from [human epithelial] ^{the conditioned medium of human fibroblast} cells as antigen. Hybridomas were prepared by routine methods and [supernatants] ^{supernatants} were screened with the ELISA assay to detect KGF-reactive antibodies. Positive clones were serially subcloned by the usual methods, and selected subclones were grown as ascites tumors in mice for production of large amounts of antibodies. Antibodies were purified from ascites fluids employing standard techniques (e.g., hydroxyapatite or immunoaffinity resins).

Polyclonal antibodies against a synthetic peptide were raised in rabbits by standard methods, as follows. The peptides were made by solid phase technology and coupled to thyroglobulin by reaction with glutaraldehyde. Serial subcutaneous injections were made and test [bleeds] ^{bleeds} were screened by ELISA as well as other techniques, including Western blot analysis and mitogenesis bioassay. IgG immunoglobulins were isolated by affinity chromatography using immobilized protein G.

Polyclonal antibodies were raised in rabbits against both naturally secreted KGF from human fibroblasts and recombinant KGF produced in *E. coli* (see next section), using the following protocol:

i) Initial injection and first boost were administered in the inguinal lymph nodes;

ii) subsequent boosts were made intramuscularly.

5

Screening of test bleeds included ELISA as well as Western blot analysis and mitogenesis bioassay, and IgG was purified as for antibodies against synthetic peptides, above.

RESULTS

Isolation of cDNA clones encoding the novel growth factor. To search for cDNA clones corresponding to the KGF coding sequence, two pools of oligonucleotides with lengths of 26 bases were generated based upon a nine-amino acid sequence, Asn-Asp-Met-Thr-Pro-Glu-Gln-Met-Ala, as determined by microsequencing of purified KGF

(see Experimental Section I, above and [reference Rabin et al., Proc. Natl. Acad. Sci. USA 86:802-806 (1989) II-3]). One oligonucleotide pool contained a mixture of all 256 possible coding sequences for the nine amino acids, while the other contained inosine residues at the degenerate third position of the codons for Thr and Pro.

This latter design reduced the number of possible coding sequences in the pool to 16.

Inosine in a tRNA anticodon can form hydrogen bonds with A, C or U (Crick, F.H.C., 1966, J. Mol. Biol. 19, 548-555 [II-4]), and oligonucleotides that contain deoxyinosine have been shown to

hybridize efficiently with the corresponding cDNA (Ohtsuka, E., Matsuki, S., Ikehara, M., Takashi, Y. and Matsubara, K., 1985, [II-5] J. Biol. Chem. 260, 2605-2608)

A cDNA library was constructed in a cDNA cloning vector, pCEV9 (Maki, T., Matsui, T., Heidaran, M., and Aaronson, S.A. [II-6] using unpublished observations.

polyadenylated RNA extracted from the human

embryonic lung fibroblast cell line M426 (Aaronson, S.A. and Todaro, G.J. [II-7] 1968, Virology 36, 254-261 the initial source of the growth factor.

Screening of the library (9×10^5 plaques) with

the ³²P-labelled 26-mer oligonucleotides identified 88 plaques which hybridized to both pools of oligonucleotide probes.

Characterization and sequencing of
5 selected cDNA clones. Of 10 plaque-purified clones that were analyzed, one, designated clone 49, had a cDNA insert of 3.5 kb, while the rest had inserts ranging from 1.8 kb to 2.1 kb.

10 Analysis of the smaller clones revealed several common restriction sites. Sequencing of a representative smaller clone, designated clone 32, along with clone 49, demonstrated that they were overlapping cDNAs (Fig⁶[II-1A]). [Whereas clone 49 was primed from the poly(A) tail of the
15 message, clone 32 arose during the construction of the library by hybridization of the oligo (dT) primer to an A-rich sequence in the 3' noncoding region of the KGF mRNA.]

Description of the sequence encoding the
20 KGF polypeptide. Alignment of the two cDNAs (clones 32 and 49) established a continuous sequence of 3.85 kb containing the complete KGF coding sequence (Fig.⁷[II-1B]). An ATG likely to be an initiation codon was located at nucleotide
25 position 446, establishing a 582-base pair open reading frame that ended at a TAA termination codon at position 1030. This open reading frame

would encode a 194-amino acid polypeptide with a calculated molecular weight of 22,512 daltons.

The sequence flanking the ATG codon did not conform to the proposed GCC(G/A)CCATGG

5 consensus for optimal initiation by eukaryotic
ribosomes ^{Kozak, M., 1987, Nucl. Acids Res. 13, 8125-8148} ((II-8)), however, there was an A three
nucleotides upstream of the ATG codon. An A at
this position is the most highly conserved
nucleotide in the consensus. This ATG codon was
10 preceded 85 nucleotides upstream by a TGA stop
codon in the same reading frame.

A 19-amino acid sequence that was homologous to the experimentally determined
[NH₂]-terminus of purified KGF began 32 amino acids
15 downstream of the proposed initiation codon.
There was complete agreement between the
predicted and experimentally determined amino
acid sequences, where unambiguous assignments
could be made.

20 To search for homology between KGF and
any known protein, a computer search of the
National Biomedical Research Foundation data base
using the FASTP program of Lipman and Pearson was
conducted ^{Lipman, D.J. and Pearson, R.W., 1985, Science 227, 1435-1443} ((II-9)). By this approach, a striking
25 degree of relatedness between the predicted
primary structure of KGF and those of acidic and

basic FGF, as well as the related ^[FGF-5] *hst* and *int-2*-encoded proteins was revealed.

Expression of mRNA transcripts of the KGF gene in human cells. In preliminary attempts to examine expression of KGF mRNA in human cells, a probe spanning the majority of the KGF coding sequence (Probe A, Figure⁶ [II-1A]) detected a single 2.4 kb transcript by Northern blot analysis of total M426 RNA (Figure⁸ [II-1D]). This was considerably shorter than the length of the composite cDNA sequence, 3.85 kb.

However, on screening poly(A)-selected M426 RNA, an additional transcript of approximately 5 kb was detected. Furthermore, a probe derived from the untranslated region of clone 49, 3' to the end of clone 32 (Probe B, Figure⁶ [II-1A]), hybridized only to the larger message (Figure⁸ [II-1C]). Thus, it appears that the KGF gene is transcribed as to alternate RNAs.

Two other members of the FGF gene family, bFGF ^{Abraham, J. A., et al., 1986, Science 233, 545-548} (II-29) and *int-2* ^{Mansour, S. L. and Martin, G. R., 1988, EMBO J. 1, 2035-2041} (II-30), also express multiple RNAs, the significance of which remains to be determined.

To investigate the normal functional role of KGF, the expression of its transcript in a variety of human cell lines and tissues was examined. As shown in Figure¹⁰ [II-3], the

predominant 2.4 kb KGF transcript was detected in each of several stromal fibroblast lines derived from epithelial tissues of embryonic, neonatal and adult sources, but not from epithelial cell lines of normal origin. The transcript was also detected in RNA extracted from normal adult kidneys and organs of the gastrointestinal tract, but not from lung or brain. The striking specificity of KGF RNA expression in stromal cells from epithelial tissues indicated that this factor plays a normal role in mesenchymal stimulation of epithelial cell growth.

For comparison, the mRNAs of other growth factors with known activity on epithelial cells were also analyzed in the same tissues as listed above. Among the epithelial and stromal cell lines analyzed, there was no consistent pattern of expression of aFGF or bFGF transcripts (Fig. ¹⁰[II-3]). The EGF transcript was not expressed [an]ⁿ any of the same cell lines, and was only observed in kidney, among the various tissues. Finally, the TGF α message was not detected in any of the stromal fibroblast lines and was expressed at varying levels in each of the epithelial cell lines. It was also detected at low levels in kidney among the tissues examined (Fig. ¹⁰[II-3]).

Inhibition of KGF mitogenic activity by

heparin. Heparin has been shown to substantially

increase the mitogenic activity of aFGF for a

variety of target cells in culture, and to

5 stabilize it from heat inactivation ([I-21, II-6138-6142 (1985), Gospodarowicz et al., Proc. Natl. Acad. Sci. USA 82: 475-485 (1986)]). Despite binding tightly to bFGF, heparin

had minimal effects on its mitogenic activity

([I-22]).^{Gospodarowicz et al., supra} In view of the relatedness of KGF to

the FGFs, the effect of heparin on KGF mitogenic

10 activity was examined. As shown in Table [II-1],³

thymidine incorporation by BALB/MK cells in

response to KGF was inhibited 16 fold when

heparin was included in the culture medium. In

contrast, the activities of both aFGF and bFGF

15 were increased by the same treatment. INSERT Table 3 from page 95.

Production of anti-KGF antibodies.

Several kinds of antibodies which recognize KGF

or KGF-like polypeptides have been prepared using

standard methodologies well known in the art of

20 experimental immunology and summarized in the

Methods section, above. These include:

monoclonal antibodies raised in mice against

intact, purified protein from human fibroblasts;

polyclonal antibodies raised in rabbits against

25 synthetic peptides with sequences based on amino

acid sequences predicted from the KGF cDNA

sequence; polyclonal antibodies raised in

rabbits against both naturally secreted KGF from

human fibroblasts and recombinant KGF produced in *E. coli* (see next section).

Monoclonal antibodies from three different hybridomas have been purified. All three recognize the recombinant as well as the naturally occurring KGF in a solid-phase (ELISA) assay. None cross-reacts with KGF under denaturing conditions (in a Western blot), and none neutralizes mitogenic activity of KGF in the BALB/MK bioassay.

Polyclonal antibodies were generated with a synthetic peptide with the amino acid sequence NDMTPEQMATNVR, corresponding to residues numbered 32 through 44 in KGF (see Fig. ⁷II-1), plus an R (^{Arg}[arg]) residue instead of the actual [asp]^{Asn} residue encoded by the cDNA at position 45. The [asp]^{Asn} residue is probably glycosylated in the natural KGF polypeptide and, therefore, [appeared to be an arg^{was not identified} in the amino acid sequencing data, appeared to be an arg].

obtained directly from that polypeptide (see Discussion, below). Polyclonal antibodies generated with this synthetic peptide recognize both naturally occurring and recombinant KGF in ELISA and Western blot analyses at a level of sensitivity of at least as low as 10 ng protein. These antibodies, however, do not neutralize mitogenic activity of KGF in the BALB/MK bioassay.

Polyclonal antisera against intact natural KGF protein recognizes KGF in both ELISA and Western blot assays. Such antibodies also appear to inhibit mitogenic activity of KGF in the BALB/MK bioassay.

Expression of KGF cDNA in *E. coli*. KGF

cDNA was expressed to produce polypeptide in *E. coli* by placing its coding sequence under control of the hybrid trk promoter (comprising elements of trp and lac promoters), in the plasmid pKK233-2 (II-31). To accomplish this, a specific length of KGF cDNA that contained the information to code for the mature KGF molecule (i.e., without

its signal peptide) was amplified using the polymerase chain reaction technique (II-32). The fragment was directionally inserted between two sites in the vector, namely the *Nco*I site, made blunt ended by *S*1 nuclease digestion, and the *Hind*III site, using standard recombinant DNA methodology. Selected recombinants were sequenced at their cDNA 5' ends to ensure correct alignment of the ATG initiation codon with the regulatory elements of the trk promoter.

Several recombinants were tested for protein production by the usual small scale methods. In brief, the clones were grown to mid-exponential phase ($OD_{595} \sim 0.5$), treated with 1 mM

Saiki, R.K., Scharf, S., Faloona, F., Mullis, K.B., Horn, G.T., Erlich, H.A. and Arnheim, N., 1985, *Science* 230, 1350-1354

isopropyl β -D-thiogalactopyranoside (IPTG) for 90 minutes, and cell extracts were run on SDS-polyacrylamide gels for Western blot analysis. All recombinants tested synthesized a protein
5 that was recognized by antibodies raised against an amino-terminal KGF peptide. One recombinant was selected which showed the greatest induction from IPTG, for further protein analyses.

One liter of bacteria was grown up in
10 NZY broth containing 50 μ g/ml ampicillin and 12.5 μ g/ml tetracycline, to OD₅₉₅ ~0.5, and treated for 90 min. with IPTG. The cells were collected by centrifugation, resuspended in 50 mM sodium phosphate (^{pH} 7.3), 0.2 M NaCl, and lysed by
15 sonication. Cell debris was removed by centrifugation, and lysate applied directly to a heparin-Sepharose affinity column.

As determined by Western blot analysis and mitogenic activity in keratinocytes,
20 recombinant KGF was eluted in 0.5-0.6 M NaCl. Subsequent purification of the HSAC material with a Mono-S (FPLC) column (Pharmacia) yielded a preparation of KGF estimated to be $\geq 90\%$ pure, as judged by electrophoretic analysis using SDS-
25 polyacrylamide gels and silver-staining.

Recombinant KGF efficiently stimulated thymidine incorporation into BALB/MK keratinocyte cells, but was only marginally active on NIH/3T3

fibroblasts. Half-maximal stimulation of the BALB/MK cells in the standard keratinocyte bioassay was achieved with a concentration of between 2 to 5 ng/ml, compared to a concentration of 10 to 15 ng/ml for KGF purified from M426 cells. One liter of bacterial cells yielded approximately 50 µg of Mono-S purified recombinant KGF.

Construction of a chimera containing KGF and aFGF sequences. The studies above indicated

that KGF possessed two distinctive characteristics which might be encoded by distinct portions or domains of the polypeptide sequence, as is well known to occur in coding sequences of other multifunctional polypeptides. To test this possibility, a chimeric DNA segment encoding the NH₂-terminal sequence of KGF grafted onto the C-terminal core of aFGF was constructed, as follows. A [Sau3AI]^[SauI] restriction enzyme site

(GATC) in the 5' end of the KGF cDNA, within codons for residues [78 and 79 (arg and ile)]^{76, 77 and 78 (Tur, Leu and Arg)} respectively; see Fig. [II-1]⁷ was cut and joined to an homologous site in the aFGF cDNA within codons for amino acids [39 (arg) and 40]^[37 (Phe), 38 (Leu) and 39 (Arg)]. The 3' and 5' ends of this chimeric DNA were joined to the vector DNA of the plasmid pKK233-2 by the same method used for insertion of the KGF cDNA

encoding the secreted form of polypeptide (see Methods, above).

When recombinant *E. coli* cells were constructed using the vector carrying the chimera, and [expressions]^{expression} tests were conducted as described for mature KGF, above, a novel product with properties of both KGF and aFGF was produced. The peptide was enriched by heparin-Sepharose chromatography and found to have a target cell preference for keratinocytes, like KGF, with minimal activity on fibroblasts (NIH/3T3). The mitogenic activity of this chimeric polypeptide lacks, however, susceptibility to inhibition by heparin, a characteristic which parallels that of aFGF rather than KGF. In fact, the mitogenic activity on keratinocytes is actually enhanced by heparin, as is the case for aFGF. Thus the peptide domains responsible for [cell target]^{target cell} specificity and heparin sensitivity are clearly distinct and readily separable in KGF, according to the practice of the present invention.

DISCUSSION

The experiments described in this section illustrate the practice of several principal embodiments of the present invention. These include isolation of cDNAs encoding KGF,

expression of such cDNAs in recombinant cells,
production of various antibodies reactive with
KGF, and construction and expression of a
chimeric cDNA encoding a novel growth factor with
5 amino acid sequences and related functionalities
of both KGF and aFGF. The following points
related to these embodiments may also be noted to
enhance the understanding of the present
invention.

10 The sequence predicted from the KGF cDNA
agreed with the amino acid sequence determined
from the purified KGF form secreted by human
fibroblasts. Moreover, the sequence offered
potential explanations for positions where
15 definitive amino acid assignments could not be
made by direct amino acid sequencing. Residues
32 and 46 are predicted from the cDNA sequence to
be cysteines, and hydrolyzed derivatives of
unmodified cysteine residues are not detectable
20 following Edman degradation. The predicted KGF
amino acid sequence also contained one potential
N-linked glycosylation site (Asn-X-Ser/Thr) from
residues 45 through 47. If Asn 45 were
glycosylated, it would not be detected by the
25 amino acid sequencing methods employed here. In
fact, KGF migrates as a broad band on
NaDodSO₄/PAGE at a higher molecular weight than

predicted for the purified protein. This may be accounted for by glycosylation.

The FGFs are heparin-binding mitogens with broad target cell specificities (II-10). ^{Thomas, K., 1987, FASEB J. 1, 434-440}

- 5 {The *hst* gene was identified as a transforming gene from a human stomach tumor (II-11), adjacent normal stomach tissue (II-12), and from Kaposi's sarcoma (II-13), by standard NIH/3T3 transfection assays. ^{Taira, et al., Proc. Natl. Acad. Sci. USA 84: 2980-2984 (1987)} ^{Yoshida, et al., Proc. Natl. Acad. Sci. USA 84: 7305-7309 (1987)} ^{Delli-Bovi, et al., Proc. Natl. Acad. Sci. USA 84: 5660-5664 (1987)} FGF-5 and *hst* are transforming genes originally detected by DNA-mediated gene transfer assays. The product of the *int-2* gene is expressed normally during mouse embryogenesis (II-14) and aberrantly after proviral integration of mouse mammary tumor virus (II-15). ^{Jakobovits, A., Shackleford, G.M., Varmus, H.E. and Martin, G.R., 1986, Proc. Natl. Acad. Sci. USA 83: 7806-7810} ^{Peters, G., Brooks, S. and Dickson, S., 1983, Cell 33, 364-377}
- 10

KGF is the sixth member of the

- fibroblast growth factor family to be identified (II-28). ^{Zhan, X., Peters, B., Hu, X. and Goldfarb, R., 1988, Mol. Cell. Biol. 8, 3487-3495} While the name FGF-6 does not seem suitable because KGF is devoid of activity on fibroblasts, this nomenclature may also be used for this growth factor, to denote its structural relationship to the FGF family. As all previously characterized growth factors either exclude epithelial cells as targets or include them among a number of sensitive target cells, the highly specific nature of KGF mitogenic activity for epithelial cells, and the sensitivity of keratinocytes in particular, make it unique.
- 15
- 20
- 25

In studies to date, expression of the KGF transcript appears to be specific for stromal cells derived from epithelial tissues, suggesting its function in normal epithelial cell proliferation. The availability of the KGF cDNA clone will make it possible to determine whether abnormal expression of this growth factor can be implicated in clinical conditions characterized by epithelial cell dysplasia and/or neoplasia. Moreover, the ability to produce large quantities of this novel growth factor by recombinant techniques should allow testing of its clinical applicability in situations where specific growth of epithelial cells is of particular importance.

Alignment of the KGF sequence with the five other proteins of the FGF family revealed two major regions of homology, spanning amino acids 65-156 and 162-189 in the predicted KGF sequence, which were separated by a short, nonhomologous series of amino acids with varying lengths in different members of the family (Fig. ⁹ [II-2]). In the case of *int-2*, the length of this sequence was 17 residues, while in *hst*, the two homologous regions were contiguous. In KGF the intervening sequence consisted of five amino acids.

In the aligned regions, the KGF amino acid sequence was about 44% identical to *int-2* (mouse), ^{41% identical to FGF-5 (human)} 39% identical to bFGF (human), 37% identical to aFGF (human) and 33% identical to *hst* (human). In this same region, all six proteins were identical at 19% of the residues, and allowing for conservative substitutions, they showed 28% homology.

As shown in Fig. ⁹[II-2], the amino termini of these related proteins are nonhomologous and of variable length. The primary KGF ^[FGF-5] and *hst* translation products contain hydrophobic N-terminal regions which likely serve as signal sequences ^{Von Heijne, G. (1986) Nucl. Acids Res. 14, 4683-4690} ([II-16]). The fact that this N-terminal domain is not present in the mature KGF molecule (Fig. [?][II-1B]) further supports this conclusion.

In contrast, the FGFs are synthesized apparently without signal peptides ^{Thomas, K., 1987, FASEB J., 1, 434-440} ([II-10]). The *int-2* protein

contains an atypically short region of N-terminal hydrophobic residues ^{Morre, R., Casati, G., Brakes, S., Dixon, M., Peters, G. and Moore, R., 1986, EMBO J. 5, 919-924} ([II-17]), but it is not known if the protein is secreted. Moreover, the *int-2* protein contains a long C-terminal extension compared to the other family members.

Purified KGF contains five cysteine residues, two of which are conserved throughout the family of FGF related proteins (Fig. ⁹[II-2]). Also of note are the five pairs of basic residues

throughout the KGF sequence. This same pattern has been observed in other FGF family members and may be involved in their interaction with heparin ^{Schwarzbauer, J.E., Tankum, J.M., Lemischka, I.R. and Hynes, R.O., 1983, Cell (II-18)}. Dibasic sites are also common targets 35, 421-431

5 for proteolytic processing and such processing might account for the microheterogeneity observed in some KGF preparations (unpublished data).

The KGF cDNA sequence was AT rich throughout its length, but particularly so in the 10 3' untranslated region where the AT content was 70% as compared to 60% in the putative coding sequence and 63% in the 5' untranslated region.

The 3' untranslated region contained a large number of ATTTA sequences, which have been 15 proposed to be involved in the selected degradation of transiently expressed, unstable RNAs ^{Shaw, G. and Kamen, R., 1986, Cell 46, 659-667 (II-19)}. There was no classical AATAAA

polyadenylation signal but two variant sequences, ^{Brnstail, M.L., Busslinger, M. and Strub, K., 1985, Cell 41, 349-359 (II-20)}, were detected 24 and 20 19 nucleotides, respectively, upstream of the poly(A) sequence at the 3' end of the cDNA.

It has been suggested that the heparin effect on acidic FGF is either due to stabilization of the active conformation of the 25 growth factor or to formation of a tertiary

complex with acidic FGF and its receptor ^{Gospodarowicz, O. and Chang, J., 1986, J. Cell Physiol. 128, 445-448 (II-21)}. If so, heparin may stabilize a

conformation of KGF that is not as active as the

^{Schreiber, A.B., Kenny, J., Kowalski, W., Friesel, J., Mehlman, T. and Maciag, i 1985, Proc. Natl. Acad. Sci. USA 82, 6138-6142}

free molecule, or form a tight complex that is
unable to efficiently interact with its receptor.

While its ability to bind heparin
reflects the structural similarities of KGF with
the FGF's, the differences in target cell
specificities between these related mitogens is
remarkable. The FGF's induce division of most
nonterminally differentiated cells of both
embryonic mesodermal and neuroectodermal origin.

In addition to fibroblasts and vascular
endothelial tissues, mesodermally derived targets

in culture include myoblasts, chondrocytes and
osteoblasts (II-23). ^{Thomas, K.A. and Gimenez-Gallego, G., 1986, Trends Biochem. Sci. 11, 86-8}
FGF's are also mitogenic
for glial astrocytes and neuroblasts (II-24). <sup>Gensburger, C., Labourdette, G.,
and Sensenbrenner, M., 1987,
FEBS Lett. 217, 1-5</sup>

The product of the oncogene isolated from
Kaposi's sarcoma, which is identical to *hst*, also

stimulates proliferation of NIH/3T3 and capillary
endothelial cells (II-25). <sup>Delli-Bovi, P., Curatola, A.A., Kern, F.G., Green, A., Itman, M,
and Fusilico, C., 1987,
Cell 50, 729-737</sup>
To date, KGF induced
mitogenesis has only been observed in epithelial

cells, and the absence of any detectable activity
in fibroblasts or endothelial cells has also been

demonstrated (see Experimental Section I, above
and (II-3)). ^{Rubin, et al., Proc. Natl. Acad. Sci. USA 86: 802-806 (1989)}
It seems likely, therefore, that KGF

acts through a different cell surface receptor
than the FGFs.

There is no significant N-terminal
homology between KGF and other FGF-related
proteins. Thus, the construction of chimeric

molecules between KGF and a prototype FGF was undertaken to determine whether the KGF N-terminal domain is sufficient to account for its unique target cell specificity. The results on the first such recombinant polypeptide sequence indicate that the N-terminal domain of KGF essentially encodes the cell preference for keratinocytes, while the susceptibility of KGF to heparin is encoded somewhere in the C-terminal core region which was replaced by sequences of aFGF. This novel KGF-like growth factor may have advantages in clinical applications where administration of an epithelial-specific growth factor is desirable in the presence of heparin, a commonly used anticoagulant. Additional studies on chimeras should also provide insights into which specific domains in the C-terminal core contribute the different effects of heparin on their biologic activities.

20

[REFERENCES FOR EXPERIMENTAL II

- II-1. James, R. and Bradshaw, R. A. (1984)
Ann. Rev. Biochem. 53, 259-292.
- II-2. Sporn, M. B. and Todaro, G. J. (1980) *N. Eng. J. Med.* 303, 878-880.
- 25 II-3. Rubin, J. S., Osada, H., Finch, P. W.,
Taylor, W. G., Rudikoff, S. and]

- [Aaronson, S. A. (1989) *Proc. Natl. Acad. Sci. USA* (in press).
- II-4. Crick, F. H. C. (1966) *J. Mol. Biol.* 19, 548-555.
- 5 II-5. Ohtsuka, E., Matsuki, S., Ikehara, M., Takashi, Y. and Matsubara, K. (1985) *J. Biol. Chem.* 260, 2605-2608.
- II-6. Miki, T., Matsui, T., Heidaran, M. and Aaronson, S. A. (1989) *Proc. Natl. Acad. Sci.* (in press).
- 10 II-7. Aaronson, S. A. and Todaro, G. J. (1968) *Virology* 36, 254-261.
- II-8. Kozak, M. (1987) *Nucl. Acids Res.* 13, 8125-8148.
- 15 II-9. Lipman, D. J. and Pearson, R. W. (1985) *Science* 227, 1435-1441.
- II-10. Thomas, K. (1987) *FASEB J.* 1, 434-440.
- II-11. Taira, M., Yoshida, T., Miyagawa, K., Sakamoto, H., Terada, M. and Sugimara, T. (1987) *Proc. Natl. Acad. Sci. USA* 84, 2980-2984.
- 20 II-12. Yoshida, T., Miyagawa, K., Odagiri, H., Sakamoto, H., Little, P. F. R., Terada, M. and Sugimara, T. (1987) *Proc. Natl. Acad. Sci. USA* 84, 7305-7309.
- 25 II-13. Delli-Bovi, P. and Basillico, C. (1987) *Proc. Natl. Acad. Sci. USA* 84, 5660-5664.]

- [II-14. Jakobovits, A., Shackleford, G. M.,
Varmus, H. E. and Martin, G. R. (1986)
Proc. Natl. Acad. Sci. USA 83, 7806-7810.
- 5 II-15. Peters, G., Brookes, S. and Dickson, S.
(1983) *Cell* 33, 364-377.
- II-16. von Heijne, G. (1986) *Nucl. Acids Res.* 14,
4683-4690.
- 10 II-17. Moore, R. Casey, G., Brookes, S., Dixon,
M., Peters, G. and Dickson, C. (1986)
EMBO J. 5, 919-924.
- II-18. Schwarzbauer, J.E., Tamkum, J.M.,
Lemischka, I. R. and Hynes, R. O. (1983)
Cell 35, 421-431.
- 15 II-19. Shaw, G. and Kamen, R. (1986) *Cell* 46,
659-667.
- II-20. Birnsteil, M. L., Busslinger, M. and
Strub, K. (1985) *Cell* 41, 349-359.
- 20 II-21. Schrieber, A. B., Kenny, J., Kowalski,
W., Friesel, J., Mehlman, T. and Maciag,
T. (1985) *Proc Natl. Acad. Sci. USA* 82,
6138-6142.
- II-22. Gospodarowicz, O. and Cheng, J. (1986)
J. Cell Physiol. 128, 475-485.
- 25 II-23. Thomas, K. A. and Gimenez-Gallego, G.
(1986) *Trends Biochem. Soc.* 11, 81-84.]

- [II-24. Gensburger, C., Labourdette, G. and Sensembrenner, M. (1987) *FEBS Lett.* 217, 1-5.
- 5 II-25. Delli-Bovi, P., Curatola, A M., Kern, F. G., Greco, A., Ittman, M. and Basilico, C. (1987) *Cell* 50, 729-737.
- II-26. Sanger, F., Nicklen, S. and Coulson, A. R. (1977) *Proc. Natl. Acad. Sci. USA* 74, 5463 5467.
- 10 II-27. Yanisch-Perron, C., Vieira, J. and Messing, J. (1985) *Gene* 33, 103-119.
- II-28. Zhan, X., Bates, B., Hu, X. and Goldfarb, M. (1988) *Mol. Cell. Biol.* 8, 3487-3495.
- 15 II-29. Abrahams J. A., Mergia, A., Whang, J.L., Tumolo, A., Friedman, J., Hjerrild, K. A., Gospodarowicz, D. and Fiddes, J. C. (1986) *Science* 233, 545-548.
- 20 II-30. Mansour, S. L. and Martin, G. R. (1988) *EMBO J.* 1, 2035-2041.
- II-31. Amman, E. and Brosius, J. (1985) *Gene* 40, 183.
- 25 II-32. Sakai, R. K., Scharf, S., Faloona, F., Mullis, K.B., Norn, G.T., Erlich, H.A. and Arnheim, N. (1985) *Science* 230, 1350-1354.]

[II-33. Beckman, P.M., Betsholtz, C. Heldin, C-
H., Westermark, B. DiMarco, E., DiFiore,
P.P., Robbins, K.C. and Aaronson, S. A.
(1988) *Science* 241, 1346-1349.]

For purposes of completing the background description and present disclosure, each of the published articles, patents and patent applications heretofore identified in this specification [are]⁵ hereby incorporated by reference into the specification.

The foregoing invention has been described in some detail for purposes of clarity and understanding. It will also be obvious that various combinations in form and detail can be made without departing from the scope of the invention.

WHAT IS CLAIMED IS:

1. A human keratinocyte growth factor (KGF) having an apparent molecular weight of about 28 kDa as determined by migration in NaDodSO₄/PAGE, and a specific activity of at least about 3.4×10^4 units per milligram of protein, where one unit of activity is defined as that amount which causes half of the maximal possible stimulation of DNA synthesis in BALB/MK keratinocyte cells under standard assay conditions.

2. Human KGF according to claim 1, wherein said specific activity is at least about 3.2×10^5 units per milligram protein.

3. A bioassay for KGF-like activity in a test sample which comprises the following steps:

- i) growing keratinocytes in culture to confluence and maintaining said confluent culture in serum-free medium;
- ii) adding a test sample to said confluent culture of keratinocytes;

and

- iii) determining the stimulation of DNA synthesis in said keratinocytes.

4. A method of producing KGF from cultured cells comprising the following steps:

- i) Culturing KGF-producing cells in culture medium under conditions such that KGF is produced;
- ii) concentrating said culture medium so that a first concentrate is formed;
- iii) contacting said concentrate with heparin under conditions such that KGF present in said first concentrate binds to the heparin whereby a heparin-KGF complex is formed;
- iv) separating said heparin-KGF complex from said concentrate;
- v) treating said heparin-KGF complex under conditions such that said KGF dissociates from the heparin so that a solution of free KGF is formed;
- vi) concentrating said solution so that a second concentrate is formed;
- vii) fractionating said second concentrate so that KGF is separated from the remaining components.

5. A method of producing KGF from cultured cells, according to claim 4, wherein said KGF-producing cells are M426 human embryonic fibroblasts.

5 6. A DNA segment encoding a human keratinocyte growth factor (KGF) protein.

7. A DNA segment, according to claim 6, wherein said protein has the amino acid sequence defined in Figure II-1.

10 8. A DNA segment encoding a chimeric KGF-like protein which comprises within a single polypeptide molecule functional segments of human KGF and at least one other polypeptide of the fibroblast growth factor family.

15 9. A recombinant DNA molecule comprising a DNA segment according to claim 6 or claim 8 and a vector.

20 10. A culture of cells transformed with said recombinant DNA molecule according to claim 9.

11. A method of producing a human KGF protein comprising culturing said cells according to claim 10 in a culture medium under conditions such that said protein is produced and isolating said protein from said cells.

12. A method of producing a human KGF protein comprising culturing said cells according to claim 10 in a culture medium, wherein said

protein is secreted from said cell, and isolating said protein from said medium.

13. A human KGF or KGF-like protein having the amino acid sequence in Figure II-1B.

5 14. A human KGF or KGF-like protein, according to claim 13, which is not glycosylated.

15. An antibody specific for a peptide having the amino acid sequence of human KGF or KGF-like protein, according to claim 13.

10 16. The antibody according to claim 15 which neutralizes the mitogenic activity of human KGF.

17. A bioassay for expression of a gene encoding KGF, comprising the steps of:.

15 i) isolating mRNA from tissues or cells;

and

ii) annealing said RNA to a DNA probe encoding a human KGF;

20 iii) determining the amount of DNA:RNA hybrid containing said DNA probe.

18. A bioassay for KGF antigen
comprising the steps of:

i) extracting polypeptides from body
fluids or tissue samples;

5 and

ii) determining the level of human KGF
antigen by reaction with an
antibody specific for a peptide
having the amino acid sequence of
human KGF or KGF like protein,
according to claim 13.

10

19. A pharmaceutical composition for
treatment of conditions requiring specific
stimulation of epithelial cells, comprising KGF
according to claim 1 or claim 13, and an
acceptable pharmaceutical carrier.

15

20. A pharmaceutical composition for
treatment of conditions requiring specific
inhibition of stimulation of epithelial cells by
KGF, comprising antibodies to KGF according to
claim 15, and an acceptable pharmaceutical
carrier.

20

ABSTRACT

Discoveries are disclosed that show particular aspects of recombinant DNA technology can be used successfully to produce hitherto
5 unknown human keratinocyte growth factor (KGF) protein free of other polypeptides. These proteins can be produced in various functional forms from spontaneously secreting cells or from DNA segments introduced into cells. These forms
10 variously enable biochemical and functional studies of this novel protein as well as production of antibodies. Means are described for determining the level of expression of genes for the KGF protein, for example, by measuring
15 mRNA levels in cells or by measuring antigen secreted in extracellular or body fluids.

and legend

Move entire Table ^{to} page 43, above line 10

Table [I-]1. Growth Factor Purification

Purification step	Protein (mg)	Total activity (units)*	Specific activity (units/mg)
Conditioned medium (10 liters)	1.4×10^{3a}	2.5×10^4	1.8×10^1
Ultrafiltration (retentate)	1.3×10^{3a}	3.2×10^4	2.5×10^1
HSAC 0.6 M NaCl pool	0.73^b	1.6×10^4	2.2×10^4
TSK-G3000 SW	8.4×10^{-3b}	2.7×10^3	3.2×10^5
C ₄ -HPLC	6.1×10^{-3b}	2.1×10^2	3.4×10^4

Recoveries were calculated by assuming that all of the mitogenic activity in the starting material was due to the isolated factor.

* One unit of activity is defined as half of the maximal stimulation of thymidine incorporation induced by TSK-purified factor in the BALB/MK bioassay, in which approximately 3 ng of the TSK-purified factor stimulated 1 unit of activity.

^a Protein was estimated by using the Bradford reagent from BioRad ([I-23]).
Bradford, M., 1976, Anal. Biochem. 72, 248-254

^b Protein was estimated by using $A_{214}^{1\%} = 140$.

move entire table to page 45, above line 24

Table [I-2]. Target Cell Specificity of Growth Factors

Growth Factor	<u>Epithelial</u>			<u>Fibroblast</u>	<u>Endothelial</u>
	BALB/MK	BS/589	CCL208	NIH/3T3S	Human saphenous vein
KGF	500-1000	2-3	5-10	<1	<1
EGF	100-200	20-40	10-30	10-20	n.d.
TGF α	150-300	n.d.	n.d.	10-20	n.d.
aFGF*	300-500	2-3	5-10	50-70	5
bFGF	100-200	2-3	2-5	50-70	5

Comparison of maximal thymidine incorporation stimulated by KGF and other growth factors in a variety of cell lines, expressed as fold stimulation over background.

- 5 This data represents a summary of four different experiments.

* Maximal stimulation by aFGF required the presence of heparin (Sigma), 20 μ g/ml.

n.d. = not determined.






Figure [I]-1. Heparin-Sepharose affinity chromatography of conditioned medium from M426 human embryonic fibroblasts. Approximately 150 ml of ultrafiltration retentate derived from five liters of M426 conditioned medium were loaded onto a heparin-Sepharose column (6 ml bed volume) in 1 hr. After washing the column with 150 ml of the equilibration buffer, 20 mM Tris-HCl, pH 7.5/0.3M NaCl, the retained protein (<5% of the total protein in the retentate) was eluted with a modified linear gradient of increasing NaCl concentration. Fraction size was 3.8 ml and flow rate during gradient elution was 108 ml/hr. Two μ l of the indicated fractions were transferred to microtiter wells containing a final volume of 0.2 ml for assay of ^3H -thymidine incorporation in BALB/MK cells as described in the Methods.

Figure [I]-2. [SEE LEGEND NEXT PAGE]




entire legend
move to p 13, line 11, after "on"


[Figure I-2. (A)] Reversed-phase C_4 HPLC of BALB/MK mitogenic activity. Active fractions eluted from heparin-Sepharose with [0-6M] ^{0.6M} NaCl were processed with the Centricon -10 and loaded directly onto a C_4 Vydac column (4.6 x 250 mm) which had been equilibrated in 0.1% trifluoroacetic acid/20% acetonitrile (ACN). After washing the column with 4 ml of equilibration buffer, the sample was eluted with a modified linear gradient of increasing % ACN. Fraction size was 0.2 ml and flow rate was 0.5 ml/min. Aliquots for the assay of 3 H-thymidine incorporation in BALB/MK cells were promptly diluted 10-fold with 50 μ g/ml bovine serum albumin/20 mM Tris-HCl, pH 7.5, and tested at a final dilution of 200-fold. (B) NaDodSO₄/PAGE analysis of selected fractions from the C_4 chromatography shown in panel A. Half of each fraction was dried, redissolved in NaDodSO₄/2-mercaptoethanol, heat denatured and electrophoresed in a 14% polyacrylamide gel which was subsequently stained with silver. The position of each molecular weight marker (mass in kDa) is indicated by an arrow. (C) DNA synthesis in BALB/MK cells triggered by the fractions analyzed in Panel B. Activity is expressed as the fold stimulation over background which was 100 cpm.



[Figure I-3.] Molecular sieving HPLC (TSK 3000SW) chromatography of the BALB/MK mitogenic activity. Approximately 50 μ l of a Centricon-processed, 0.6M NaCl pool from HSAC were loaded onto a LKB GlasPac TSK G3000SW column (8 x 300 mm), previously equilibrated in 20 mM Tris-HCl, pH 6.8/0.5M NaCl, and eluted as 0.2 ml fractions at a flow rate of 0.4 ml/min. Aliquots of 2 μ l were transferred to microtiter wells containing a final volume of 0.2 ml for assay of ³H-thymidine incorporation in BALB/MK cells. The elution positions of molecular weight markers (mass in kDa) were as indicated by the arrows.



[Figure I-4.] Comparison of BALB/MK DNA synthesis in response to TSK-purified mitogen and other growth factors. Incorporation of ^3H -thymidine into trichloroacetic acid-insoluble DNA, expressed as fold stimulation over background, was measured as a function of the concentration of the indicated growth factors. Background values with no sample added were 150 cpm. The results represent mean values of two independent experiments. Replicates in each experiment were within 10% of mean values. TSK-purified mitogen, $\text{---}\bullet\text{---}$; EGF, $\text{---}\triangle\text{---}$; aFGF, $\text{---}\square\text{---}$; bFGF, $\text{---}\circ\text{---}$.



[Figure I-5.] Comparative growth of BALB/MK cells in a chemically defined medium in response to different combinations of growth factors.

Cultures were plated at a density of 2.5×10^4

cells per dish on 35 mm Petri dishes precoated with poly-D-lysine/fibronectin in a 1:1 mixture of Eagle's minimal essential medium and Ham's F12 medium supplemented with transferrin,

Na_2SeO_3 , ethanolamine and the growth factors

indicated below. After 10 days, the plates were fixed and stained with Giemsa. Key: a) no

growth factor; b) EGF alone; c) insulin alone;

d) KGF alone; [e) EGF + insulin] e) EGF and dialyzed fetal calf serum (final concentration, 10%); f) KGF and EGF; g) KGF and insulin; h) EGF and insulin

concentrations of the growth factors were as

follows: EGF, 20 ng/ml; insulin, 10 $\mu\text{g/ml}$; and

KGF, 40 ng/ml.

move to p. 14, line 6, after "shows"

and legend
move entire table to p. 63, line 15,
after "treatment."

3
TABLE [II-1]. Effect of Heparin on KGF Mitogenic Activity.

Growth Factor	BALB/MK		NIH/3T3	
	[+] -	[-] +	[+] -	[-] +
KGF	150	9.5	<1	<1
aFGF	106	259	10.4	68
bFGF	30	124	45.7	70

Cells were plated in microtiter plates, grown to confluence in serum containing media and then placed in a serum-free medium for 24-72 hr prior to sample addition. Mitogenesis assays were performed as described (see Experimental Section I, above and [II-3]). Where indicated, heparin was included in the culture media at a final concentration of 20 μ g/ml. The concentration of all the growth factors was 50 ng/ml. The results represent fold stimulation of 3 H-thymidine incorporation in the indicated assay cell in the presence (+) or absence (-) of heparin. Each value represents the mean result from two independent experiments in which each point, in turn, represents the mean value of duplicate analyses.

Rubin et al. Proc. Natl. Acad. Sci. USA 86: 802-806 (1989)

Figure [II-1A. SEE LEGEND FOLLOWING] 6

A.





Figure II-1B. SEE LEGEND FOLLOWING

၈၂

100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918

Figure [II-1C. SEE LEGEND FOLLOWING] 8


C.



move to page 14, line 10, before "Table"


[Figure II-1. Nucleotide sequence and deduced amino acid sequence of KGF cDNA, and identification of KGF gene transcripts. (A)] Figure 6 outlines a schematic representation of human KGF cDNA clones. Overlapping pCEV9 clones 32 and 49, used in sequence determination, are shown above a diagram of the complete structure in which untranslated regions are depicted by a line and the coding sequence is boxed. The hatched region denotes sequences of the signal peptide [and the open region of the mature protein]. Selected restriction sites are indicated. [(B)] Figure 7 documents the KGF cDNA nucleotide and predicted amino acid sequences. Nucleotides are numbered on the [right]; ^{left} amino acids are numbered throughout. The N-terminal peptide sequence derived from purified KGF is underlined. The hydrophobic N-terminal domain is italicized. The potential asparagine-linked glycosylation site is overlined. The variant polyadenylation signals, AATTAA and AATACA, close to the 3' end of the RNA, are boxed. [(C)] ^{Figure 8 shows} Identification of KGF mRNAs by Northern blot analysis. Lanes a and c, poly(A)-selected M426 RNA; [lane d], ^{lanes b and d} total cellular M426 RNA. Filters were hybridized with a ^h ^{32}P -labeled 695 bp *Bam*HI/*Bcl*I fragment from clone 32 (Probe A, Fig. [II-1A]), lanes a and b,

or a 541 bp *ApaI/EcoRI* fragment from clone 49
(Probe B, Fig. ⁶II-1A), lanes c and d.
1



5 [Figure II-2. Topological comparison of the FGF family of related molecules. The two protein domains that share high homology are shown by shaded boxes. Hatched boxes indicate putative signal peptide sequences. The positions of two conserved cysteine residues (C) are shown.]

Figure [II-3. SEE LEGEND NEXT PAGE] 10



entire legend
move to page 15, line 10, before "fig."

10 shows

Figure [II-3.] Northern blot analysis of KGF mRNA in normal human cell lines and tissues, and comparison with mRNA expression of other growth factors with known activity on epithelial cells.

5 Total cellular RNAs were isolated by cesium trifluoro-acetate gradient centrifugation. 10 μ g of RNA were denatured and electrophoresed in 1% formaldehyde gels. Following mild alkali denaturation (50 mM NaOH for 30'), RNA was transferred to nitrocellulose filters using 1 M ammonium acetate as a convectant. Filters were hybridized to a 32 P-labelled cDNA probe

containing the [647bp EcoRI fragment from the 5' end] of the KGF coding sequence (A) or similar

647 bp EcoRI fragment from the 5' end

15 probes from the other growth factor DNAs. The following human cell types were used: squamous cell carcinomas (A253, A388 and A431); mammary epithelial cells (B5/S89); immortalized branchial epithelial cells (S6 and R1); keratinocytes immortalized with Ad12-SV40; primary human
20 keratinocytes; neonatal foreskin fibroblasts, (AG1523); adult skin fibroblasts (501T); and embryonic lung fibroblasts (WI-38 and M426).